

**Nutrient Broth****M002**

Nutrient Broth is used for the general cultivation of less fastidious microorganisms, can be enriched with blood or other biological fluids.

**Composition\*\***

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Sodium chloride	5.000
Beef extract	1.500
Yeast extract	1.500
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

**Directions**

Suspend 13 grams in 1000 ml distilled water. Heat, if necessary, to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

**Principle And Interpretation**

Nutrient media are basic culture media used for maintaining microorganisms, cultivating fastidious organisms by enriching with serum or blood and are also used for purity checking prior to biochemical or serological testing (1, 2). Nutrient Broth has the formula originally designed for use in the Standard Method for Examination of Water and Waste water. It is one of the several non-selective media useful in routine cultivation of microorganisms (3, 4). It can be used for the cultivation and enumeration of bacteria which are not particularly fastidious. Addition of different biological fluids such as horse or sheep blood, serum, egg yolk etc. makes it suitable for the cultivation of related fastidious organisms. Peptic digest of animal tissue, beef extract and yeast extract provide the necessary nitrogen compounds, carbon, vitamins and also some trace ingredients necessary for the growth of bacteria. Sodium chloride maintains the osmotic equilibrium of the medium.

**Quality Control****Appearance**

Cream to yellow homogeneous free flowing powder

**Colour and Clarity of prepared medium**

Light amber coloured clear solution in tubes

**Reaction**

Reaction of 1.3% w/v aqueous solution at 25°C. pH : 7.4±0.2

**Cultural Response**

M002: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth				
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant				
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good-luxuriant				
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good-luxuriant				
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	good-luxuriant				

**Reference**

- Lapage S., Shelton J. and Mitchell T., 1970, Methods in Microbiology', Norris J. and Ribbons D., (Eds.), Vol. 3A, Academic Press, London.
- MacFaddin J. F., 2000, Biochemical Tests for Identification of Medical Bacteria, 3rd Ed., Lippincott, Williams and Wilkins, Baltimore.
- Downes F. P. and Ito K., (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.
- American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.

**Storage and Shelf Life**

Store below 30°C and the prepared medium at 2 - 8°C. Use before expiry date on the label.