



Tryptone Soya-Tryptose Broth

M1876

Intended Use:

Recommended for identification of *Salmonella* species from food samples in accordance with FDA BAM, 1998.

Composition**

Ingredients	Gms / Litre
Tryptone	8.500
Soya peptone	1.500
Sodium chloride	5.100
Dextrose (Glucose)	1.770
Dipotassium hydrogen phosphate	1.250
Tryptose	10.380
Yeast extract	3.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 31.50 grams in 1000 ml purified / distilled water. Heat if necessary to dissolve the medium completely. Dispense 5ml portions into 16×150mm test tube. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Salmonella is a genus of rod-shaped, Gram-negative, non-spore-forming, predominantly motile enterobacteria with peritrichous flagella. Most of the species are pathogenic, and the infections are mainly due to the ingestion of contaminated food (1). Tryptone Soya Tryptose Broth is used for serological identification of *Salmonella* species with respect to 'polyvalent flagellar (H) test' in accordance with FDA BAM, 1998 (2).

Add 25g of the food sample(s) suspected to be contaminated with *Salmonella* into 225ml culture broth (1:9 ratio) and incubate at $35 \pm 2.0^\circ \text{C}$ for 24 ± 2.0 hours in accordance with the BAM protocol. The incubated sample is processed for isolation of the species by inoculation into selective media such as Selenite broth (M052), Fluid Tetrathionate Medium w/o Iodine and BG, Modified (M032F) or Rappaport Vassiliadis Medium, Modified (M880F) and incubation for 24hrs at appropriate temperatures. Thoroughly mix and streak a 3 mm loopful of the incubated broth on Bismuth Sulphite Agar (M027), XLD agar (M031F), and Hektoen Enteric Agar, w/ 1.2% agar (M467F). Organism is identified by its colony characteristics in respective media. The organism can be confirmed through biochemical and serological tests. Serological tests include identification of polyvalent flagellar (H) antigen. Tryptone soya tryptose broth (M1876) is used for the initial inoculum preparation for this test (2,3).

Tryptone, Tryptose, Papaic digest of soyabean meal and Yeast extract provide necessary nitrogen compounds and vitamin source and Dextrose provide necessary carbon source to the medium. Sodium chloride maintains the osmotic equilibrium of the medium. Dipotassium hydrogen phosphate acts as the buffering agent.

Type of specimen

Food samples

Specimen Collection and Handling:

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,5,7). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions:

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations:

1. Further biochemical and serological tests must be carried out for further identification.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution without any precipitate.

Reaction

pH of 3.15% w/v aqueous solution at 25°C . pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth
<i>Staphylococcus aureus</i> ATCC 25923 (00034*)	50 -100	luxuriant
<i>Escherichia coli</i> ATCC 8739 (00012*)	50 -100	luxuriant
<i>Escherichia coli</i> ATCC 25922 (00013*)	50 -100	luxuriant
<i>Pseudomonas aeruginosa</i> ATCC 27853 (00025*)	50 -100	luxuriant
<i>Salmonella Typhi</i> ATCC 6539	50 -100	luxuriant
<i>Salmonella Enteritidis</i> ATCC 13076 (00030*)	50 -100	luxuriant
<i>Salmonella Typhimurium</i> ATCC 14028	50 -100	luxuriant
<i>Salmonella Abony</i> NCTC 6017	50 -100	luxuriant

Key : *Corresponding WDCM numbers.

Reference

1. MacFaddin, J. F. 1985. Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria vol. 1. Baltimore: Williams and Wilkins.
2. FDA, U.S. 1998. Bacteriological Analytical Manual. 8 ed. Gaithersburg, MD: AOAC International.
3. Forbes, B. A., Sahm, D. F. and Weissfield, A. S. 2002. Bailey and Scott's Diagnostic Microbiology. 11 ed. St Louis: The C.V. Mosby Co.

Revision : 0 / 2012

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.