



Tergitol-7 Agar Base

M616

Tergitol-7 Agar Base is recommended for selective enumeration and identification coliform organisms.

Composition**

Ingredients	Gms / Litre
Proteose peptone	5.000
Yeast extract	3.000
Lactose	10.000
Sodium heptadecyl sulphate(Tergitol 7)	0.100
Bromo thymol blue	0.025
Agar	15.000
Final pH (at 25°C)	6.9±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 33.12 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add 3 ml of Triphenyl Tetrazolium Chloride (TTC) Solution (FD057), if desired. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Tergitol-7 Agar was originally designed by Chapman (1) and later on modified by incorporating 2,3,5-Triphenyl Tetrazolium Chloride (TTC) into the medium. This medium is selective and differential used for the detection and enumeration of coliform organisms. Pollard (2) has reported the selective bactericidal property of sodium heptadecyl sulphate (Tergitol-7). Kulp et al (3) corroborated the use of Tergitol-7 Agar with TTC in routine analysis of water and Mossel (4) used this medium for the examination of food materials.

Proteose peptone and yeast extract serve as sources of carbon, nitrogen and other essential nutrients including vitamin B complex. Sodium heptadecyl sulphate (Tergitol-7) inhibits gram-positive bacteria and *Proteus* swarming and yields better recovery of coliforms. Bromo thymol blue is the pH indicator. Lactose fermenting organisms form yellow colonies with yellow zones while *Klebsiella* and *Enterobacter* form greenish yellow colonies. Lactose non-fermenters produce blue colonies. TTC is reduced by the bacterial cell except *Escherichia coli* and *Enterobacter aerogenes* to form formazan, a red coloured insoluble complex, thereby producing red coloured colonies.

Filter the specimen to be analyzed through two membranes. Place the membrane upon two TTC Tergitol Agar plates. Incubate one plate at 37°C for 24 hours (total coliforms) and the other at 44°C for 18-24 hours (faecal coliforms). The yellow colonies with deep yellow halo after incubation at 44°C should be identified as faecal coliform bacteria.

Quality Control

Appearance

Cream to light green homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Green coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.31% w/v aqueous solution at 25°C. pH : 6.9±0.2

pH

6.70-7.10

Cultural Response

M616: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours with added TTC Solution 1% (FD057).

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony/ medium
Cultural Response				
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	>=50%	Reddish brown
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	>=50%	yellow with red centre
<i>Proteus mirabilis</i> ATCC 25933	50-100	good	40-50%	red with bluish zone
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good	40-50%	red with bluish zone
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	>=50%	red with bluish zone
<i>Staphylococcus aureus</i> ATCC 25923	>=10 ³	inhibited	0%	
<i>Shigella flexneri</i> ATCC 12022	50-100	good-luxuriant	>=50%	red with bluish zone

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

- 1.Chapman G.H., 1947, J. Bact., 53:504.
- 2.Pollard A.L., 1946, Science, 103:758.
- 3.Kulp W., Mascoli C. and Tavshanjian O., 1953, Am. J. Public Health, 43:1111.
- 4.Mossel D.A.A., 1962, J. Appl. Bact., 25:20.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.