



Soyabean HiVeg Medium, Sterile powder

MV011G

Soyabean HiVeg Medium, Sterile powder is prepared by completely replacing animal based peptones with vegetable peptones. It is a gamma irradiated sterile powder recommended for the evaluation of sterility in manufacturing process.

Composition**

Ingredients	Gms / Litre
HiVeg hydrolysate	17.000
Papaic digest of Soyabean meal	3.000
Sodium chloride	5.000
Dipotassium phosphate	2.500
Dextrose	2.500
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Sterile powder can be used directly for the evaluation of sterility in manufacturing process. For sterile liquid medium aseptically add 30 grams in 1000 ml sterile distilled / purified water. Heat if necessary to dissolve the medium completely. DO NOT AUTOCLAVE OR OVERHEAT. Excessive heating is detrimental. Dispense aseptically in sterile tubes or flasks as desired.

Principle And Interpretation

Soyabean HiVeg Medium, Sterile powder is prepared by completely replacing animal based Pancreatic digest of casein of Soyabean Casein Digest Medium, Sterile powder with vegetable based HiVeg hydrolysate. This makes the medium free of BSE/ TSE associated risks. This medium can be used for the same purpose of Soyabean Casein Digest Medium, which is recommended by various pharmacopoeias as sterility testing medium (1, 2). It is also used for the sensitivity testing by the tube dilution method for antimicrobial agents (3).

The combination of HiVeg hydrolysate and papaic digest of soyabean meal makes this medium nutritious by providing amino acids and long chain peptides for the growth of microorganisms. Dextrose serves as the carbon source and dipotassium phosphate buffers the medium. Sodium chloride maintains the osmotic balance of the medium.

Quality Control

Appearance

Cream to yellow may have green tinge homogeneous free flowing powder

Colour and Clarity of prepared medium

Light amber coloured clear solution

Reaction

Reaction of 3.0% w/v aqueous solution at 25°C. pH : 7.3±0.2

pH

7.10-7.50

Sterility Testing

No growth is observed after 14 days for Bacteria at 30-35°C and for Fungi at 20-25°C.

Test for Mycoplasma (PCR)

None detected.

Stability test

Light yellow coloured clear solution without any precipitation or sedimentation at room temperature for 7 days

Growth promoting properties

Clearly visible growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating ≤ 100 cfu (at 30-35°C for 18-24 hours).

Sterility Testing + Validation

The medium is tested with suitable strains of microorganisms inoculating ≤ 100 cfu and incubating at 20-25°C for not more than 3 days in case of bacteria and not more than 5 days in case of fungi.

Cultural Response

Organism	Inoculum (CFU)	Growth	Incubation temperature	Incubation period
Growth promoting				
<i>Staphylococcus aureus</i> ATCC 6538	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Staphylococcus aureus</i> ATCC 25923	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Escherichia coli</i> ATCC 8739	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Escherichia coli</i> ATCC 25922	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Escherichia coli</i> NCTC 9002	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Pseudomonas aeruginosa</i> ATCC 9027	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Pseudomonas aeruginosa</i> ATCC 27853	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Bacillus subtilis</i> ATCC 6633	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Micrococcus luteus</i> ATCC 9341	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Salmonella Typhimurium</i> ATCC 14028	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Salmonella Abony</i> NCTC 6017	50 -100	luxuriant	30 -35 °C	18 -24 hrs
<i>Streptococcus pneumoniae</i> ATCC 6305	50 -100	luxuriant	30 -35 °C	18 -24 hrs
Sterility Testing- Growth promotion+ Validation				
<i>Staphylococcus aureus</i> ATCC 6538	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Staphylococcus aureus</i> ATCC 25923	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Escherichia coli</i> ATCC 8739	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Escherichia coli</i> ATCC 25922	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Escherichia coli</i> NCTC 9002	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Pseudomonas aeruginosa</i> ATCC 9027	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Pseudomonas aeruginosa</i> ATCC 27853	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Bacillus subtilis</i> ATCC 6633	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Micrococcus luteus</i> ATCC 9341	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Salmonella Typhimurium</i> ATCC 14028	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Salmonella Abony</i> NCTC 6017	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Streptococcus pneumoniae</i> ATCC 6305	50 -100	luxuriant	20 -25 °C	≤ 3 d
<i>Candida albicans</i> ATCC 10231	50 -100	luxuriant	20 -25 °C	≤ 5 d
<i>Candida albicans</i> ATCC 2091	50 -100	luxuriant	20 -25 °C	≤ 5 d
* <i>Aspergillus brasiliensis</i> ATCC 16404	50 -100	luxuriant	20 -25 °C	≤ 5 d

Storage and Shelf Life

Store below 10-30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

Reference

- 1.The United States Pharmacopoeia / National Formulary, 2012, 35.The United States Pharmacopoeial Convention Inc.,Rockville, MD.
- 2.Indian Pharmacopoeia, 2014, Govt. of India, Ministry of Health and Family Welfare, New Delhi, India.
- 3.Wright and Welch, 1959-60, Antibiotics Ann., 61.

Revision : 02 / 2015



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.