

Pseudomonas Agar (For Pyocyanin), Granulated

GM119

Pseudomonas Agar (For Pyocyanin), granulated is recommended for the detection of pyocyanin production by *Pseudomonas* species.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	20.000
Potassium sulphate	10.000
Magnesium chloride	1.400
Agar	15.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 46.4 grams in 1000 ml distilled water containing 10 ml glycerol. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Pseudomonas Agar is based on the formulation described by King et al (1) and as recommended in U.S. Pharmacopoeia (2) for detecting pyocyanin, a water soluble pigment by *Pseudomonas* species (3). This medium enhances the elaboration of pyocyanin but inhibits the formation of fluorescein pigment. The fluorescein pigment diffuses from the colonies of *Pseudomonas* into the agar and shows blue colouration. Some *Pseudomonas* strains produce small amounts of fluorescein resulting in a blue-green colouration.

Potassium sulphate and magnesium chloride, which enhances the pyocyanin production and suppresses the fluorescein production. A pyocyanin-producing *Pseudomonas* strain will usually also produce fluorescein. It must, therefore, be differentiated from other simple fluorescent pseudomonads by other means. Temperature can be a determining factor as most other fluorescent strains will not grow at 35°C. Rather, they grow at 25-30°C (3).

Quality Control

Appearance

Cream to yellow coloured granular medium

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.64% w/v aqueous solution containing 1% v/v glycerol at 25°C. pH:7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural response was observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Medium
<i>Pseudomonas aeruginosa</i> ATCC 9027	50 -100	luxuriant	>=50 %	blue-green
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	luxuriant	>=50 %	blue-green

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

- 1.King, Ward and Raney, 1954, J.Lab. and Clin. Med., 44:301
- 2.The United States Pharmacopoeia, 2014, The United States Pharmacopoeial Convention, Rockville, MD.
- 3.MacFaddin J., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.