

## Xylose-Lysine-Deoxycholate Agar, Granulated

GMH031

Xylose-Lysine Deoxycholate Agar, granulated is a selective medium recommended for the isolation and enumeration of *Salmonella* Typhi and other *Salmonella* species from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP.

### Composition\*\*

Ingredients	Gms / Litre
Xylose	3.500
L-lysine	5.000
Lactose monohydrate	7.500
Sucrose	7.500
Sodium chloride	5.000
Yeast extract	3.000
Sodium deoxycholate	2.500
Sodium thiosulphate	6.800
Ferric ammonium citrate	0.800
Phenol red	0.080
Agar	13.500
pH after heating ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 54.8 grams (the equivalent weight of dehydrated medium per litre) in 1000 ml purified/ distilled water. Heat with frequent agitation until the medium boils. DO NOT HEAT IN AN AUTOCLAVE. Transfer immediately to a water bath at 50°C. After cooling, pour into sterile Petri plates. It is advisable not to prepare large volumes, which will require prolonged heating and may produce precipitate.

Note: Slight precipitation in the medium may occur, which is inherent property of the medium, and does not affect the performance of the medium.

### Principle And Interpretation

*Enterobacteriaceae* is a family of gram-negative, non-spore-forming bacilli that contains more than 100 species of bacteria that normally inhabit the intestines of humans and animals. Members forming part of the normal intestinal microflora are referred to as coliforms. The clinically significant genera of *Enterobacteriaceae* include *Cedecea*, *Citrobacter*, *Edwardsiella*, *Enterobacter*, *Escherichia*, *Ewingella*, *Hafnia*, *Klebsiella*, *Kluyvera*, *Proteus*, *Salmonella*, *Shigella* and *Yersinia* (1).

The Salmonellae are the most complex of all the *Enterobacteriaceae*. Human *Salmonella* infections are most commonly caused by ingestion of food, water or milk, contaminated by human or animal excreta (2). A large number of media have been developed for the selective isolation and identification of enteric bacilli including *Salmonella*.

Xylose Lysine Deoxycholate Agar is a selective as well as differential medium formulated by Taylor (3-7) for the isolation and identification of enteric pathogens especially Shigellae from stool samples. It is also used for pharmaceutical testing and non-sterile product testing for the detection (or absence) of *Salmonella* after enrichment in Rappaport Vassiliadis Salmonella Enrichment Broth (MH1491/GMH1491) in accordance with the harmonized method of USP/EP/BP/JP/IP (8-12).

Deoxycholate, ferric ammonium citrate and sodium thiosulphate are selective agents that inhibit gram-positive microorganisms. Essential nutrients, growth factors for growth of microorganism are provided by yeast extract. Xylose, sucrose and lactose are the fermentable sugars in this medium. Xylose is fermented by almost all the enteric bacteria except Shigellae, which enable the differentiation of Shigellae from Salmonellae. Salmonellae metabolize the xylose and decarboxylate lysine and thus change the pH to alkaline and mimic Shigellae reaction. However to prevent this reaction by lysine positive coliforms, lactose and sucrose are added in excess to produce acid and hence nonpathogenic H<sub>2</sub>S producers do not decarboxylate lysine. Sodium thiosulphate helps in reactivation of sulphur containing compounds and prevents the desiccation of these compounds during storage. It also forms the substrate for enzyme thiosulphate reductase, which breaks it to form H<sub>2</sub>S. Thiosulphate and ferric

ammonium citrate are the H<sub>2</sub>S indicators in the medium. Sodium thiosulphate is also inactivator of halogens, mercurial and aldehyde and can minimize its toxicity in the testing sample, if any during microbial limit tests. Sodium chloride maintains the osmotic equilibrium in this medium. Phenol red is the pH indicator.

Degradation of fermentable sugars proceed concurrently and generates acids, which cause pH indicator to give various shades of colour, causing a color change in the colonies and in the medium from red to yellow on prolonged incubation. Hydrogen sulfide production results in colonies with black centers under alkaline conditions, which can be inhibited by acid production by carbohydrate fermentation. Alkaline condition causes the color of the medium to change back to red. This medium is an ideal medium for screening samples containing mixed flora of enteric pathogens as recovery of Salmonellae and Shigellae is not conspicuous by even profuse growth of other species (13,14).

## Quality Control

### Appearance

Light yellow to pink coloured granular medium

### Gelling

Firm, comparable with 1.35% Agar gel

### Colour and Clarity of prepared medium

Red coloured clear to slightly opalescent gel forms in Petri plates

### pH

7.20-7.60

### Cultural Response

Growth Promotion is carried out in accordance with the harmonized method of USP/EP/BP/JP/IP. Cultural response was observed after an incubation at 30-35°C for specified time. Recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar.

### Growth promoting properties

Growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating ≤100 cfu(at 30-35°C for ≤18 hours).

### Indicative properties

Colonies are comparable in appearance and indication reaction to those previously obtained with previously tested and approved lot of medium occurs for the specified temperature for a period of time within the range specified inoculating ≤100cfu (at 30-35°C for 18-72 hours).

### Inhibitory properties

No growth of the test microorganism occurs for the specified temp for not less than longest period of time specified inoculating ≥100cfu (at 30-35°C for ≥ 72 hours).

### Cultural Response

Cultural characteristics observed after incubation at 30-35 °C for 18-48 hours. Recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony	Incubation temperature
<b>Growth Promoting + Indicative</b>					
<i>Salmonella</i> Typhimurium ATCC 14028	50 -100	luxuriant	≥50 %	red with black centres	18 -72 hrs
<i>Salmonella</i> Abony NCTC 6017	50 -100	good-luxuriant	≥50 %	red with black centres	18 -72 hrs
<b>Additional Microbiological testing</b>					
<i>Escherichia coli</i> ATCC 8739	50 -100	fair	20 -30 %	yellow	18 -72 hrs
<i>Escherichia coli</i> ATCC 25922	50 -100	fair	20 -30 %	yellow	18 -72 hrs
<i>Escherichia coli</i> NCTC 9002	50 -100	fair	20 -30 %	yellow	18 -72 hrs
<i>Proteus vulgaris</i> ATCC 13315	50 -100	good-luxuriant	≥50 %	grey with black centres	18 -72 hrs
<i>Salmonella</i> Paratyphi A ATCC 9150	50 -100	good-luxuriant	≥50 %	red	18 -72 hrs

<i>Salmonella</i> Paratyphi B ATCC 8759	50 -100	good-luxuriant	$\geq 50$ %	red with black centres	18 -72 hrs
<i>Salmonella</i> Enteritidis ATCC 13076	50 -100	good-luxuriant	$\geq 50$ %	red with black centres	18 -72 hrs
<i>Salmonella</i> Typhi ATCC 6539	50 -100	good-luxuriant	$\geq 50$ %	red with black centres	18 -72 hrs
<i>Shigella dysenteriae</i> ATCC 13313	50 -100	good-luxuriant	$\geq 50$ %	red	18 -72 hrs
<i>Shigella flexneri</i> ATCC 12002	50 -100	fair-good	30 -40 %	red	18 -72 hrs
<i>Shigella sonnei</i> ATCC 25931	50 -100	fair-good	30 -40 %	red	18 -72 hrs
<i>Enterobacter aerogenes</i> ATCC 13048	50 -100	fair	20 -30 %	yellow	18 -72 hrs
<i>Enterobacter cloacae</i> ATCC 13047	50 -100	fair	20 -30 %	yellow	18 -72 hrs
<i>Staphylococcus aureus</i> ATCC 25923	$\geq 10^3$	inhibited	0%		$\geq 72$ hrs
<i>Staphylococcus aureus</i> ATCC 6538	$\geq 10^3$	inhibited	0%		$\geq 72$ hrs
<i>Enterococcus faecalis</i> ATCC 29212	$\geq 10^3$	inhibited	0%		$\geq 72$ hrs

## Storage and Shelf Life

Store below 30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry date on the label.

## Reference

1. Pelczar M. J. Jr., Reid R. D., Chan E. C. S., 1977, Microbiology, 4th Ed., Tata McGraw-Hill Publishing Company Ltd, New Delhi.
2. Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company.
3. Taylor W. L., 1965, Am. J. Clin. Pathol., 44:471-475.
4. Taylor W. L. and Harris B., 1965, Am. J. Clin. Pathol., 44:476.
5. Taylor W. L. and Harris B., 1967, Am. J. Clin. Pathol., 48:350.
6. Taylor W. L. and Schelhart B., 1967, Am. J. Clin. Pathol., 48:356.
7. Taylor W. L. and Schelhart B., 1968, Am. J. Clin. Pathol., 16:1387.2.
8. The United States Pharmacopoeia, 2014, The United States Pharmacopoeial Convention, Rockville, MD.
9. British Pharmacopoeia, 2014, The Stationery Office British Pharmacopoeia.
10. European Pharmacopoeia, 2014, European Department for the Quality of Medicines of Council of Europe.
11. Japanese Pharmacopoeia, 2008, Published by Society of Japanese Pharmacopoeia, Tokyo, Japan.
12. Indian Pharmacopoeia, 2014 Ministry of Health and Family Welfare, Govt. of India.
13. McCarthy M.D., 1966, N.Z. J. Med. Lab. Technol., 20:127.
14. Isenberg H.D., Kominos S. and Siegal M., 1969, Appl. Microbiol., 18:656.

Revision : 00 / 2014



### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.