



Diluting Fluid A

LQ121L

Diluent in testing of pharmaceuticals in accordance with USP.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	1.000
Final pH (at 25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Diluting fluid is used as the diluting or rinsing solution for membrane filter techniques in pharmaceutical products. Measured portions of Diluting fluid A should be used to rinse the membrane after filtration. Inoculate this rinse with 50-100 cfu of test organisms. Simultaneously run a positive control of the same medium. Incubate both the set of medium at the specified time and temperature. Compare the growth obtained for the rinse with that obtained in the positive control after incubation.

Principle And Interpretation

Diluting Fluid A is recommended as rinsing fluid for membrane filter method used in validation tests for bacteriostasis and fungistasis activity of pharmaceutical articles before carrying out sterility test procedures as per USP (1). After filtering the specified quantity of the test specimen, the membrane is rinsed with measured portions of rinsing or diluting fluid. This rinse is inoculated with known number of test bacteria and fungi as specified in pharmacopoeia. The resultant growth is compared with positive control to determine presence of fungistasis or bacteriostasis activity in test specimen.

Quality Control

Appearance

Sterile clear Diluting Fluid A in bottle.

Colour

Light yellow coloured medium

Quantity of medium

300ml of medium in bottle

pH

6.90-7.30

Sterility test

Passes release criteria.

Growth Promotion Test

In accordance with the harmonized method of USP.

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours

Organism	Growth	Inoculum (CFU)
<i>Candida albicans</i> ATCC 10231	good	50-100
<i>Escherichia coli</i> ATCC 25922	good	50-100
<i>Staphylococcus aureus</i> ATCC 25923	good	50-100
<i>Staphylococcus aureus</i> ATCC 6538	good	50-100

Storage and Shelf Life

Store between 2-8°C. Use before expiry date on the label.

Reference

1. The United States Pharmacopoeia / National Formulary, USP34 / NF29, 2011, US Pharmacopeial convention Inc., Rockville, MD.



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.