



## Buffered Sodium Chloride Peptone Solution

LQ123L

For the preparation of test suspension in accordance with harmonized methods of USP,EP & JP

### Composition\*\*

Ingredients	Gms / Litre
Potassium dihydrogen phosphate	3.600
Disodium hydrogen phosphate dihydrate	7.200
Sodium chloride	4.300
Peptone (meat or casein)	1.000

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Label the ready to use LQ123L bottle. Inoculate the sample and Incubate at specified temperature and time.

### Principle And Interpretation

The composition of this medium is as per USP (1) and is in accordance with the harmonized methodology of USP/EP/BP/JP (1,2,3,4). This medium is recommended for preparation of stable test strain suspension employed for validating the microbiological testing procedures of non-sterile products. The standardized stable suspensions are used so that the suitability of this test to detect microorganism in presence of product can be established. Non-fatty products insoluble in water and water-soluble products are diluted/dissolved using this solution. Peptone (meat or casein) serves as nutrient source and maintains the cell viability. Phosphates in the medium act as good buffering agents. Sodium chloride maintains the osmotic balance and cell integrity. Polysorbates reduce surface tension and also inactivate phenolic compound, if present in the test sample. Edel and Kampelmacher (6) noted that sub lethal injury to Salmonellae might occur in many food preservation processes. Pre-enrichment in Buffered Sodium chloride-Peptone solution pH 7.0 (LQ123L) at 35°C for 18-24 hours results in repair of injured cells (5). This medium supports the repair of injured cells that have sensitivity to low pH. It is also recommended for pre-enrichment and repair of injured cells (5).

### Quality Control

#### Appearance

Sterile clear Buffered Sodium Chloride Peptone Solution in bottle.

#### Colour

Colourless solution.

#### Quantity of medium

300ml of medium in bottle

#### pH

7.00- 7.00

#### Growth Promotion Test

In accordance with the harmonized method of USP/EP/BP/JP.

#### Sterility test

Passes release criteria

#### Cultural response

Cultural characteristics observed after an incubation at 30-35°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth
<i>Staphylococcus aureus</i> ATCC25923	50 -100	luxuriant
<i>Escherichia coli</i> ATCC 25922	50 -100	luxuriant
<i>Escherichia coli</i> NCTC 9002	50 -100	luxuriant
<i>Salmonella</i> Abony NCTC 6017	50 -100	luxuriant
<i>Salmonella</i> Typhimurium ATCC 14028	50 -100	luxuriant
<i>Bacillus subtilis</i> ATCC 6633	50 -100	luxuriant
<i>Candida albicans</i> ATCC10231	50 -100	luxuriant
<i>Candida albicans</i> ATCC 2091	50 -100	luxuriant
<i>Staphylococcus aureus</i> ATCC 6538	50 -100	luxuriant
<i>Pseudomonas aeruginosa</i> ATCC 27853	50 -100	luxuriant
<i>Pseudomonas aeruginosa</i> ATCC 9027	50 -100	luxuriant
<i>Escherichia coli</i> ATCC 8739	50 -100	luxuriant

## Storage and Shelf Life

Store between 2-8°C. Use before expiry date on the label.

## Reference

1.The United States Pharmacopoeia, 2011, The United States Pharmacopoeial Convention. Rockville, MD. 2.British Pharmacopoeia, 2011, The Stationery office British Pharmacopoeia 3.European Pharmacopoeia, 2011, European Dept. for the quality of Medicines. 4.Japanese Pharmacopoeia, 2008. 5.Sadovski A.Y., 1977, J. Fd. Technol., 12:85. 6.Edel W. & Kampelmacher E.H.,1973, Bull, Wld. Hlth. Org., 48:167.

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

