



Broth Medium D (Lactose Monohydrate Broth)

M1003B

Lactose monohydrate Broth is used for detection of coliform bacteria in water, foods, dairy products in accordance with British Pharmacopoeia.

Composition**

Ingredients	Gms / Litre
Pancreatic digest of gelatin	5.000
Beef extract	3.000
Lactose monohydrate	5.000
pH after sterilization	6.9±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 12.75 grams of dehydrated powder in 1000 ml purified/distilled water. Heat if necessary to dissolve the medium completely. For larger inocula (10 ml or more), concentrated medium may be prepared to account for medium dilution by the inoculum. Dispense in tubes containing inverted fermentation vial (Durhams tube) as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Lactose Broth is recommended by British pharmacopoeia(4) for selective pre-enrichment of *Enterobacteriaceae* as well as for *E.coli* and *Salmonella* in water, food and pharmaceutical products. The medium is used for detection of specified microorganisms of non-sterile products according to British Pharmacopoeia. This medium is also recommended by various other pharmacopoeia (5,6,7)

Lactose monohydrate broth is recommended by APHA in the performance and confirmation of the presumptive test for coliform bacteria in water (1), food (2) and milk (3). This medium can be used as an alternate to Buffered sodium chloride-peptone solution pH 7.00 for the revival of *Enterobacteriaceae* and gram negative bacteria. The medium is incubated for a time sufficient to revive the bacteria but not the multiplication of the bacteria. It is recommended to incubate the medium usually for 2 ours and not for more than 5 hours.(4)

Pancreatic digest of gelatin and beef extract supply essential nutrients to the organisms. Lactose monohydrate is a fermentable carbohydrate.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light amber coloured clear solution without any precipitate

pH of 1.27% w/v aqueous solution after sterilization.

pH

6.70-7.10

Growth Promotion Test

Cultural characteristics observed after an incubation at 35-37°C for specified time

Cultural Response

Organism	Inoculum (CFU)	Recovery within 2 hours of incubation	Recovery within <=5 hours of incubation
----------	----------------	---------------------------------------	---

Preparation of test strain

<i>Escherichia coli</i> ATCC 8739	50 -100	no increase in colony count	no increase in colony count
<i>Pseudomonas aeruginosa</i>	50 -100	no increase in colony count	no increase in colony count
ATCC 9027			

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2- 8°C. Use before expiry date on the label.

Reference

- 1.Eaton A. D., Clesceri L. S. and Greenberg A W.,(Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st ed., APHA, Washington, D.C.
- 2.Downes F P and Ito K(Eds.), 2001, Compendium of Methods For The Microbiological Examination of Foods, 4th ed., APHA, Washington, D.C
- 3.Wehr H M and Frank J H., 2004, Standard Methods for the Examination of Dairy Products, 17th ed., APHA Inc., Washington, D.C.
- 4.British Pharmacopoeia, 2009, The Stationery office British Pharmacopoeia.
- 5.The United States Pharmacopoeia, 2008, The United States Pharmacopoeial Convention. Rockville, MD.
- 6.European Pharmacopoeia, 2008, European Department, for the quality of Medicines.
- 7.The Indian Pharmacopoeia 2007, Govt. of India, 2007. The Controller of Publication, Delhi.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.