



M-Tryptone Glucose Extract Broth

M1116

M-Tryptone Glucose Extract Broth is used for enumeration of bacteria in milk and other samples of sanitary importance in dairy industries by membrane filter technique.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Dextrose	2.000
Beef extract	6.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 18 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

The dairy industry has relied for safety on control of the manufacturing process and on the use of tests such as the standard plate count and coliform count as indicators of post process contamination. Testing of dairy products or dairy plant environment samples for pathogens has not been routinely performed. However, there is a need for at least surveillance testing of product and environmental samples as well as for re-evaluation of processing and environmental control procedures. Evaluation of this may require that product and environmental samples be analyzed for pathogens (4).

M-Tryptone Glucose Extract Broth is used as non-selective general purpose media recommended by APHA (1) for determination of bacterial counts in dairy products and water (2), foods (3) and other specimens respectively.

Casein enzymic hydrolysate and beef extract provide the essential nutrients like amino acids, minerals and trace growth factors. Dextrose serves as the carbon source. About 2 ml of the broth medium is used to saturate sterile absorbent pads. Filters used for membrane filtration are then aseptically placed on these absorbent pads.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light amber coloured clear solution without any precipitate

Reaction

Reaction of 1.8% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

M1116: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth
Cultural Response		
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant
<i>Salmonella Typhi</i> ATCC 6539	50-100	luxuriant
<i>Staphylococcus aureus</i> ATCC 25923	50-100	luxuriant

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

- 1.American Public Health Association, 1960, Standard Methods for the Examination of Water and Wastewater, 11th Ed., APHA, New York.
- 2.Greenberg A. E., Trussell R. R. and Clesceri L. S. (Eds.), 1985, Standard Methods for the Examination of Water and Wastewater, 16th Ed., APHA, Washington, D.C.
- 3.Speck M. L., (Ed.), 1984, Compendium of Methods for the Microbiological Examination of Foods, 2nd Ed., APHA, Washington, D.C.
- 4.American Public Health Association, 1992, Standard Methods for the Examination of the Dairy Products, 16th Ed., APHA, Washington, D.C.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.