



Carbohydrate Consumption Broth Base

M1264

Intended Use:

Recommended for cultivation and differentiation of *Listeria* species.

Composition**

Ingredients	Gms / Litre
Proteose peptone	10.000
Sodium chloride	5.000
HM peptone B #	1.000
Bromocresol purple	0.100
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

- Equivalent to Beef extract

Directions

Suspend 16.1 grams in 990 ml purified / distilled water. Heat if necessary to dissolve the medium completely. Dispense into tubes containing inverted Durhams tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Aseptically add 10 ml separately sterilized carbohydrate solution to give a final concentration of 0.5%. Mix well.

Principle And Interpretation

Carbohydrate Consumption Broth is used for the cultivation and differentiation of *Listeria* species and formulated as per Atlas (1). It is also recommended by FDA (2) and ISO (3) with a slight difference in the concentration of bromocresol purple. Differentiation is based on fermentation of glucose, xylose, rhamnose, ribose, a-methyl-D-mannoside and mannitol. Proteose peptone and HM peptone B in the medium provide carbon and nitrogen compounds including essential amino acids, vitamins and trace ingredients for bacterial metabolism. Bromocresol purple is the pH indicator, which indicates acid production by turning yellow in colour.

Carbohydrate utilization test: Inoculate each kind of carbohydrate fermentation broth with one loopful of inoculum. Incubate for 7 days at 37°C. Observe daily for acid induced colour change and gas formation. Sometimes weak positive reactions may occur after 48 hours of incubation (2).

Type of specimen

Pure isolates from food samples.

Specimen Collection and Handling

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,2,8).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Limitations:

NA

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to beige homogeneous free flowing powder

Colour and Clarity of prepared medium

Purple coloured, clear solution without any precipitate

Reaction

Reaction of 1.61% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH

6.60-7.00

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	w/o carbohydrate acid	w/o carbohydrate gas	w/ rhamnose (acid)	w/ rhamnose (gas)
Cultural Response <i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	negative reaction, no colour change	negative reaction	positive reaction, yellow colour	positive reaction
<i>Listeria monocytogenes</i> ATCC 19111	50-100	good-luxuriant	negative reaction, no colour change	negative reaction	positive reaction, yellow colour	negative reaction
<i>Listeria monocytogenes</i> ATCC 19112	50-100	good-luxuriant	negative reaction, no colour change	negative reaction	positive reaction, yellow colour	negative reaction
<i>Listeria monocytogenes</i> ATCC 19117	50-100	good-luxuriant	negative reaction, no colour change	negative reaction	positive reaction, yellow colour	negative reaction
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good-luxuriant	negative reaction, no colour change	negative reaction	negative reaction, no colour change	negative reaction
<i>Listeria monocytogenes</i> ATCC 19118	50-100	good-luxuriant	negative reaction, no colour change	negative reaction	positive reaction, yellow colour	negative reaction

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

1. Atlas R. M., 2004, Handbook of Microbiological Media, 3rd Edition, CRC Press, Washington D. C.
2. FDA Bacteriological Analytical Manual, 2005, 18th Ed., AOAC, Washington, D.C.
3. International Organization for Standardization (ISO), 1993, Draft ISO/DIS 10560.

Revision : 02 / 2016

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.