



Buffered Peptone Water w/ NaCl

M1275

Intended use

Recommended as a diluent for carrying microbial limit test from clinical and non clinical specimens.

Composition**

Ingredients	Gms / Litre
Peptone	1.000
Potassium dihydrogen phosphate	3.560
Disodium hydrogen phosphate	7.230
Sodium chloride	4.300
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 16.09 grams in 1000 ml purified / distilled water. Heat if necessary to dissolve the medium completely. Add 0.1 to 1% w/v polysorbate 20 or 80 if desired. Dispense in tube or flasks as desired. and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Buffered Peptone Water is a pre-enrichment medium designed to help recovery of sub-lethally damaged *Salmonellae* before transfer to a selective medium. This pre-enrichment medium is free from inhibitors and is well buffered and provides conditions for resuscitation of the cells that have been injured by processes of food preservation. It was noted by Edel and Kampelmacher (4) that sub-lethal injury to *Salmonella* may occur due to food preservation techniques involving heat, desiccation, high osmotic pressure, preservatives or pH changes. Buffered Peptone Water during the pre-enrichment period helps in recovery of injured cells that may be sensitive to low pH (10). This is particularly important for vegetable specimens, which have low buffering capacity. These media can be used for testing dry poultry feed (9). In a survey involving isolation of *Salmonellae* from meat that had been artificially contaminated with sub-lethally injured organisms. Pre-enrichment in Buffered Peptone Water at 37°C for 18 hours before selection in Tetrathionate Brilliant Green Bile Broth (M1255) showed superior results compared with direct selection method.

Lactose Broth is frequently used as a pre-enrichment medium but it may be detrimental to recovery of *Salmonellae* (2). The composition of Buffered Peptone Water with NaCl medium is as per IP and EP specifications recommended to dilute the sample for microbial examination (6, 5). Depending on the amount of fat in the sample to examine the kind and quantity of emulsifying agent to be used.

These pre-enrichment media contain peptone as a source of carbon, nitrogen, vitamins and minerals. Sodium chloride maintains the osmotic balance and phosphates buffer the medium. The broth is rich in nutrients and produces high resuscitation rates for sublethally injured bacteria and supports intense growth. The phosphate buffer system prevents bacterial damage due to changes in the pH of the medium. Inoculate 10 grams specimen in 50 ml of these media and incubate at 35-37° C for 18 hours. Transfer 10 ml from this medium to 100 ml of Tetrathionate Broth (M032) and incubate at 43°C for 24-48 hours and then subculture on selective plating media. Examine the plates for characteristic *Salmonella* colonies .

Type of specimen

Clinical samples - feces sample samples; Food and dairy samples

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (7,8).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,3,11). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

Limitations :

This medium contains low nutrients and hence is not recommended for the growth of organisms.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

White to cream homogeneous free flowing powder

Colour and Clarity of prepared medium

Colourless to pale yellow clear solution without any precipitate

pH

6.80-7.20

Cultural response

Cultural characteristics observed after recovery on Soybean Casein Digest Agar after an incubation at 30-35°C for 18-24 hours for bacteria and Sabouraud Dextrose Agar at 30-35°C for 24-48 hours .

Organism	Inoculum (CFU)	Recovery within 2 hours of incubation	Recovery within 4 hours of incubation	Recovery within 24 hours of incubation
<i>Escherichia coli</i> ATCC 8739 (00012*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Escherichia coli</i> ATCC 25922 (00013*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Escherichia coli</i> NCTC 9002	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Staphylococcus aureus</i> ATCC 6538 (00032*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Staphylococcus aureus</i> ATCC 25923 (00034*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Pseudomonas aeruginosa</i> ATCC 9027 (00026*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Pseudomonas aeruginosa</i> ATCC 27853 (00025*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Salmonella Typhimurium</i> ATCC 14028 (00031*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)

<i>Salmonella Abony</i> NCTC 6017 (00029*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Bacillus subtilis</i> subsp. <i>spizizenii</i> ATCC 6633 (00003*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Micrococcus luteus</i> ATCC 9341	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Candida albicans</i> ATCC 10231	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Candida albicans</i> ATCC 2091	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)

* - Corresponding WDCM Numbers

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (7,8).

Reference

1. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
2. Angelotti, 1963, Academic Press, New York, N.Y.
3. Downes F. P. and Ito K., (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.
4. Edel and Kampelmacher, 1973, Bull. W.H.O., 48:167.
5. European Pharmacopoeia, 2008, European Directorate For The Quality of Medicine
6. Indian Pharmacopoeia, 1997, Ministry of Health and Family Welfare, Govt. of India, Vol. 2.
7. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
8. Jorgensen, J.H., Pfäller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
9. Juven, Cox, Bailey, Thomson, Charles and Schutze, 1984, J. Food Prot., 47:299.
10. Sadvoski, 1977, J. Food Technol., 12:85.
11. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
