



Neutralizing Fluid

M1420B

This medium is recommended for neutralizing the activity of antimicrobial agents in accordance with BP.

Composition**

Ingredients	Gms / Litre
Peptone (meat or casein)	1.000
Lecithin (egg)	3.000
Histidine hydrochloride	1.000
Sodium chloride	4.300
Potassium dihydrogen phosphate	3.600
Disodium hydrogen phosphate, dihydrate	7.200

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 18.64 grams (the equivalent weight of dehydrated medium per litre) in 1000 ml distilled water containing 30 gm of polysorbate 80. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15lbs pressure (121°C) for minutes.

Principle And Interpretation

Neutralising fluid is used to neutralize the activity of antimicrobial agents generally present in pharmaceutical materials(1). This is required to neutralize the effect of antimicrobials while testing the sterility of such materials. This medium may be added to Buffered Sodium Chloride Peptone Solution, pH 7.0 before sterilization. If utilized their efficacy and non-toxicity towards microorganisms are demonstrated (1).

The neutralising agents present in the medium neutralises the activity of antimicrobial agents present in various pharmaceutical products which may interfere with microbial limit tests or sterility testing analysis. Egg lecithin and polysorbate 80 act as neutralising agents. Sodium chloride maintains osmotic equilibrium and phosphates serve as buffering agents.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Colourless to pale yellow opalescent solution in tubes

Cultural Response

Cultural characteristics observed when subcultured on Tryptone Soya Agar (M290), after an incubation at 35-37°C for 40-48 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth
Cultural Response		
<i>Bacillus subtilis</i> ATCC 6633	50-100	good
<i>Escherichia coli</i> ATCC 8739	50-100	good
<i>Pseudomonas aeruginosa</i> ATCC 9027	50-100	good
<i>Staphylococcus aureus</i> ATCC 6538	50-100	good
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good

Please refer disclaimer Overleaf.

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. British Pharmacopoeia, 2009, The Stationery office British Pharmacopoeia.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.