

Buffered Peptone Water

M1494I

Hmsdmcde trd

Buffered Peptone Water is used as pre-enrichment medium for increasing the recovery of injured *Salmonella* species from foods prior to selective enrichment and isolation. The composition and performance criteria of this medium are as per the applications laid down in ISO 6579-2002.

Composition**

	Gms / Litre
Hmfqdchdmsr	
Tryptone	10.000
Sodium chloride	5.000
Disodium hydrogen phosphate.12H2O	9.000
Potassium dihydrogen phosphate	1.500
FinalpH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters
Dpthu kdms sn Dmyxl shb chfdrs ne b rdhm

Directions

Suspend 20.07 grams(equivalent weight of dehydrated medium) in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Microorganisms that are subjected to environmental stresses may become structurally or metabolically damaged or injured. These microorganisms are unable to replicate in selective environments. Therefore these injured organisms must be resuscitated or permitted to repair the damage by incubation in an appropriate, non-selective environment (6). Edel and Kampelmacher (2) noted that sublethal injury to Salmonellae may occur in many food preservation processes. Enriching injured cells in Lactose Broth (pH 6.9) may be further detrimental to their recovery (0). Pre-enrichment in Buffered Peptone Water (M1494I) at 35°C for 18-24 hours results in repair of injured cells (5). The buffering system prevents bacterial damage due to change in the pH of the medium. Recently ISO committee has also recommended this pre-enrichment medium for the detection of *Enterobacteriaceae* from food stuffs and other materials (2).

Inoculate 10 grams specimen in 50 ml of Buffered Peptone Water (M1494I) and incubate at 35°C for 18 hours. Transfer 10 ml from this medium to 100 ml of Mueller Kauffman Tetrathionate Novobiocin Broth Base (M1496I) and Rappaport Vassiliadis Soya Broth (RVS Broth) (M1491) and incubate at 43°C for 24-48 hours and then subculture on selective media like XLD Agar, Modified (M031I). Examine the plates for colonies of *Salmonella* species.

6 [R G Q H U R G E K O G P

Food and dairy samples

Specimen Collection and Handling

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (3). After use, contaminated materials must be sterilized by autoclaving before discarding.

qmhmf mc Oqdb tshnmr

d c sgd k adkadenqd nodmhsd bnms hmdq-d q oqnsdbshudknudr.oqnsdbshudknsgmf.d dqnsdbshnmr bd oqnsdbshnm- Enkknv fnnc lhbqnahknfnhb k k a oq bshbdr vghkd g mckhmf rodbhldmr mc btkstqd- Rs mc qc oqdb tshnmr r odq drs akhrgcd fthcdkhmdr rgnk c ad enkknvdc vghkd g mckhmfrodbhldmr- R eds fthcdkhmdr l ad qdedqqdchm hmchuhct k r eds c s rgddsr

Khllhs shnmr

Ctd sn mtsqshnm k u qh shnmr rnl d rsq hmr l rgnv onnq fqnvsg-

Odqenql mbd mc Du kt shnm

Odqenql mbd ne sgd ldchtl hr dwodbsdc vgd m trdc r odq sgd chqdbshnm nm sgd k adk vhsghdwdgd odqhnc vgd m rsnqdc s qdbnlldmcde sdldq stqd-

Please refer disclaimer Overleaf.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution without any precipitate

Reaction

Reaction of 2.0% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.(Recovery is observed on XLD Agar, M0311)

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response			
<i>Salmonella Enteritidis</i> ATCC 50-100 13076 (00030*)		luxuriant	≥50%
<i>Salmonella Typhi</i> ATCC 6539	50-100	luxuriant	≥50%
<i>Salmonella Typhimurium</i> ATCC 14028 (00031*)	50-100	luxuriant	≥50%
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	fair-good	30-40%
<i>Pseudomonas aeruginosa</i> ATCC 27853 (00025*)	50-100	luxuriant	≥50%

Key : * Corresponding WDCM numbers

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Reference

1. Angelotti R., 1963, "Microbiological Quality of Foods", Academic Press, New York.
2. Edel W. and Kampelmacher E. H., 1973, Bull. Wld. Hlth. Org., 48: 167.
3. International Organization for Standardization (ISO), 2017, Draft ISO/DIS, 6579.
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
6. Sadovskii A. Y., 1977, J. Food Technol., 12.85.
7. Vanderzant C. and Splittstoesser D. F., (Eds.), 1992, Compendium of Methods for the Microbiological Examination of Foods, 3rd Ed., APHA, Washington, D.C.

Revision : 04/2018

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.