



Burkholderia Cepacia Agar Base

M1640

Burkholderia Cepacia Agar Base is a selective medium used for isolation of *Burkholderia cepacia* from the respiratory secretions of patients with cystic fibrosis and other non-clinical specimens

Composition**

Ingredients	Gms / Litre
Peptone	5.000
Yeast extract	4.000
Sodium pyruvate	7.000
Potassium dihydrogen phosphate	4.400
Disodium hydrogen phosphate	1.400
Bile salts	1.500
Ammonium sulphate	1.000
Magnesium sulphate	0.200
Ammonium ferrous sulphate	0.010
Phenol red	0.020
Crystal violet	0.001
Agar	12.000
Final pH (at 25°C)	6.2±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 18.26 grams in 500 ml purified / distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 50°C and aseptically add the rehydrated contents of 1 vial of Burkholderia Selective Supplement (FD232). Mix well and pour in sterile Petri plates.

Principle And Interpretation

Burkholderia cepacia is an important opportunistic pathogen and causes pulmonary infection among individuals with cystic fibrosis (CF). The organism may lead to *Burkholderia cepacia* syndrome, a neutralizing pneumonia associated with fever that culminates in to a rapid and fatal clinical deterioration (7). *B. cepacia* is difficult to isolate on routinely used laboratory media like MacConkey Agar, since *B. cepacia* is a slow grower and therefore it is usually outgrown by the faster growing *Escherichia coli*, *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Burkholderia Cepacia Agar is based on PC medium, which was originally devised by Gilligan (2). This medium was found to be superior to MacConkey Agar for growth of *B. cepacia*. The medium is made selective for *B. cepacia* by the incorporation of bile salts, crystal violet and antibiotics. The antibiotics included are Polymyxin B, Gentamycin, Ticarcillin in the form of freeze dried supplement (FD). Peptone and yeast extract in the medium provides the nitrogenous, vitamin B source and other essential nutrients. Crystal violet, bile salts and antimicrobial agents are used as selective agents. Crystal violet and bile salts inhibits gram-positive cocci including Enterococci and Staphylococci. The antibiotics (FD) namely ticarcillin, polymyxin B and gentamycin inhibit gram-negative bacteria. *B. cepacia* metabolizes pyruvate forming alkaline end products. These end products elevate the pH of the medium. The phenol red indicator changes colour from pink orange to pink red in alkaline pH.

Inoculate the plate with the specimen so as to obtain isolated colonies. The plates should be incubated for a period of 4 days to allow *B. cepacia* to grow and form colonies and subsequent colour change (2,6). The medium is not selective only for *B. cepacia*. Other organisms forming similar colonies may also grow on this medium. Therefore results obtained on this media should not be the sole criteria for identification of *B. cepacia* (1).

Type of specimen

Clinical samples - Throat, Sterile body fluids

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (4,5). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use only. Read the label. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

Limitations :

1. Some strains may show poor growth due to varying nutritional requirements.
2. Further biochemical tests must be carried out for confirmation.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 1.2% Agar gel.

Colour and Clarity of prepared medium

Orange coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.65% w/v aqueous solution at 25°C. pH : 6.2±0.2

pH

6.00-6.40

Cultural Response

Cultural characteristics observed, with addition of Burkholderia Selective Supplement (FD232), after an incubation at 35-37°C for 48-72 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Burkholderia cepacia</i> ATCC 25608	50-100	good-luxuriant	≥50%	sage green colonies with bright pink medium
<i>Pseudomonas aeruginosa</i> ATCC 9027 (00026*)	≥10 ⁴	inhibited	0%	

Key : * - Corresponding WDCM numbers

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Reference

1. Christensen et al, 1980, J. Clin. Microbiol., 27:27
2. Gilligan, 1996. Clin. Microbiol. Newsl. 18:8
3. Gilligar, Gage, Bradshaw, schidlow and Deciscco, 1985, J. Clin. Microbiol., 22
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W.(2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
6. MacDonald Gilligan, Welch, Reller and Menegus, 1994, Vol. 5:1, Cystic Fibrosis Foundation, Washington, D.
7. Whitby P. W., 1998, J. Clin. Microbiol., 36:1642 1645


Revision : 02 / 2019



In vitro diagnostic medical device



CE Marking



Storage temperature



} v^{uP}
uP



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg,Mumbai-86,MS,India



CE Partner 4U ,Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.