



A-7 Agar Base (Shepard's Differential Agar Base)

M1739

Intended Use

This medium is used for the cultivation and differentiation of *Ureaplasma urealyticum* from urine, based on its ability to produce ammonia from urea. Also used for the cultivation of other *Ureaplasma* species.

Composition**

Ingredients	Gms / Litre
Tryptone	12.830
Agar	15.000
Sodium chloride	3.800
Soya peptone	2.280
Dipotassium hydrogen phosphate	1.900
Dextrose (Glucose)	1.900
Manganese sulphate	0.730
Yeast extract	2.440
Final pH (at 25°C)	6.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 40.88 grams in 790 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add 195 ml sterile Horse serum (RM1239), 1 vial of Urea solution (FD253) (Filter sterilized) and rehydrated contents of 1 vial each of Ureaplasma Selective Supplement (FD254) and Ureaplasma Growth Supplement (FD255). Mix well and dispense as desired.

Principle And Interpretation

This medium is used for selective cultivation of urogenital *Mycoplasma*, viz. *Ureaplasma urealyticum* from clinical samples based on its ability to produce ammonia from urea (1). The medium contains tryptone and soya peptone which provide nitrogen and carbon compounds, long chain amino acids, vitamins and other necessary nutrients for the growth of *Ureaplasma*. Yeast extract provides preformed nucleic acid precursors, necessary for the growth of fastidious *Ureaplasma*. Glucose is the carbohydrate source. Sodium chloride maintains the osmotic balance.

Many *Mycoplasma* / *Ureaplasma* require serum for their good growth and also the presence of antibiotics is necessary to prevent the growth of contaminating organisms. *Ureaplasma* possess the enzyme urease and hence breakdown the urea to ammonia. Bacteria that produce ammonia appear as golden to dark brown coloured colonies (1).

Type of specimen

Clinical samples - Urine

Specimen Collection and Handling:

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (2,3).

Warning and Precautions :

In Vitro diagnostic Use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

Limitations:

1. Although this medium is selective for gram negative organisms, biochemical identification and serological testing using pure cultures is recommended for complete identification.
2. It is advised to incubate for recommended period and temperature to avoid misinterpretation of results.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of Prepared medium

Light yellow coloured clear to slight opalescent solution forms in Petri plates.

Reaction

Reaction of 4.08% w/v aqueous solution at 25°C. pH : 6.0±0.2

pH

5.80-6.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours, with added RM1239, FD253, FD254, FD255.

Cultural Response

Organism	Inoculum (CFU)	Growth	Urea
Cultural Response <i>Ureaplasma urealyticum</i> ATCC 14027	50-100	Good-luxuriant	positive reaction,

Storage and Shelf Life

Store below 30°C in a tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

Reference

1. Atlas R. M. 2004, Handbook of Microbiological Media, 3rd Ed, CRC Press, Boca Raton, Florida.
2. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.


Revision : 02/ 2017



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg, Mumbai-86, MS, India



CE Partner 4U, Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.