



Aleksandrow Broth

M1997

For enrichment of Potassium solubilizing bacteria from soil samples.

Composition**

Ingredients	Gms / Litre
Magnesium sulphate	0.500
Calcium carbonate	0.100
Potassium alumino silicate	2.000
Glucose	5.000
Ferric chloride	0.005
Calcium phosphate	2.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 9.60 grams in 1000ml of distilled water. Heat if necessary to dissolve the medium completely. Dispense into tubes or flasks as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Soil potassium supplementation relies heavily on the use of chemical fertilizer, which has a considerable negative impact on the environment. Potassium-solubilizing bacteria convert insoluble potassium in the soil into a form that plants can access. A wide range of bacteria namely *Pseudomonas*, *Burkholderia*, *Acidithiobacillus ferrooxidans*, *Bacillus mucilaginosus*, *Bacillus edaphicus*, *B.circulans* and *Paenibacillus* sp. has been reported to release potassium in accessible form from potassium-bearing minerals in soils (1). Potassium-solubilizing bacteria have been reported to exert beneficial effects on growth of cotton, pepper and cucumber(2), sorghum(3), wheat(4) and Sudan grass(5). Therefore potassium solubilizing bacteria are extensively used as biofertilizers.

Salts present in the medium support the growth of potassium solubilizing bacteria by providing the essential nutrients. The source of potassium salts is potassium alumino silicates. Potassium solubilizing bacteria will grow on this medium and form a clear zone around the colony, formed due to potassium solubilization in the vicinity of the colony.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Cream to light yellow coloured solution with white precipitate forms in tubes

Reaction

Reaction of 0.96% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth
Cultural Response <i>Potassium solubilizing isolate</i>	50-100	good-luxuriant

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Sheng, X.F., 2005, Growth promotion and increased potassium uptake of cotton and rape by a potassium releasing strain of *Bacillus edaphicus*. *Soil Biology and Biochemistry*, 37, 1918-1922
2. Han, H.S., Supanjani and Lee, K.D., 2006, Effect of co-inoculation with phosphate and potassium solubilizing bacteria on mineral uptake and growth of pepper and cucumber. *Plant Soil and Environment*, 52, 130-136.
3. Badr, M .A., Shafei, A.M . and Sharaf, S.H. El-Deen, 2006, The dissolution of K and phosphorus bearing minerals by silicate dissolving bacteria and their effect on sorghum growth. *Research Journal of Agriculture and Biological Sciences*, 2, 5-11.
4. Sheng, X.F. and He, L.Y., 2006, Solubilization of potassium bearing minerals by a wild type strain of *Bacillus edaphicus* and its mutants and increased potassium uptake by wheat. *Canadian Journal of Microbiology*, 52, 66-72.
5. Basak, B.B. and Biswas, D.R., 2008, Influence of potassium solubilizing microorganism (*Bacillus mucilaginosus*) and waste mica on potassium uptake dynamics by sudan grass(*Sorghum vulgare Pers*) grown under two Alfisols. *Plant Soil*, 317, 235-255

Revision : 00 / 2015



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.