



GN Broth, Hajna

M242

GN Broth, Hajna, is recommended for selective enrichment of gram-negative enteric organisms.

Composition**

Ingredients	Gms / Litre
Tryptose	20.000
Dextrose	1.000
Mannitol	2.000
Sodium citrate	5.000
Sodium deoxycholate	0.500
Dipotassium phosphate	4.000
Monopotassium phosphate	1.500
Sodium chloride	5.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 39 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense in test tubes. Sterilize by autoclaving at 115°C (10 lbs pressure) for 15 minutes. AVOID EXCESSIVE HEATING.

Principle And Interpretation

Hajna (1) developed Gram Negative (GN) Broth as an enrichment medium for recovery of *Salmonella* and *Shigella* from clinical and non-clinical specimens such as urine, blood clots, throat swabs, swabs from eating and drinking utensils etc (1, 2). GN Broth, Hajna is also recommended by APHA (3) for the microbiological examination of foods. Croft and Miller isolated more strains of *Shigella* from rectal swabs using this medium (9). Taylor and Schelhart showed the superiority of GN Broth to selenite enrichment media for isolation of *Shigella* (10). Hajna (2, 4) also suggested the enrichment of organisms from rectal swabs in this medium 1-6 hours before plating on solid media.

The medium contains tryptose, which provides amino acids and other nitrogenous substances to support bacterial growth. The combination of sodium citrate and sodium deoxycholate inhibit gram-positive and some gram-negative bacteria such as coliforms. Phosphates serve as a buffering system. Sodium chloride maintains osmotic equilibrium. The higher concentration of mannitol over dextrose limits the growth of *Proteus* and enhances growth of mannitol fermenting *Salmonella* and *Shigella*. *Proteus*, *Pseudomonas* and coliforms do not overgrow *Salmonella* and *Shigella* in GN Broth during the first 6 hours of incubation. This enrichment broth should be used in conjunction with selective and nonselective plating media to increase the probability of isolating pathogens (3, 5, 7).

GN Broth, Hajna should be inoculated directly with the specimen. In case of stool specimens, approximately 1 gram should be used for inoculation. Appropriate references for processing of clinical and food samples should be followed (3, 5, 6, 8). After incubation of 6-8 hours and again after 24 hours, sub culturing on selective agar media should be carried out (7).

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light amber coloured, clear to slightly opalescent solution in tubes.

Reaction

Reaction of 3.9% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

M242: Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.

Organism	Inoculum (CFU)	Growth in GN broth	Growth after 24 hours on MacConkey Agar	Colour of colony
Cultural Response				
<i>Escherichia coli</i> ATCC 25922	50-100	good	good	pink-red with bile ppt
<i>Enterococcus faecalis</i> ATCC 19433	50-100	none-poor	none-poor	pale pink-red
<i>Proteus mirabilis</i> ATCC 25933	50-100	good	good	colourless
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good	good	colourless
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good	good	colourless
<i>Shigella flexneri</i> ATCC 12022	50-100	good	good	colourless

Storage and Shelf Life

Store below 30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

Reference

- Hajna A. A., 1955, Publ. Health Lab., 13:59.
- Hajna A. A., 1955, Publ. Health Lab., 13:83.
- Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.
- Hajna A. A., 1956, Air. Univ. Sch. Ar. Med., USAF, 56:39
- Murray P. R., Baron J. H., Pfaller M. A., Tenover J. C. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
- Forbes B. A., Sahm A. S., and Weissfeld D. F., Bailey & Scotts Diagnostic Microbiology, 10th Ed., 1998, Mosby, Inc., St. Louis, Mo.
- MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
- Ewing, 1986, Edwards and Ewings Identification of Enterobacteriaceae, 4th Ed., Elsevier Science Publishing Co., Inc., New York, N.Y.
- Croft C. C., Miller M. J., 1956, Am. J. Clin. Pathol., 26:411.
- Taylor W.I., Schelhart D., 1968, Appl. Environ. Microbiol., 16:1383.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.