



0c p p k v q n " U c n v " D t q v j

05:5

K p v g p f g f " W u g <

Recommended for selective isolation of presumptive pathogenic Staphylococci.

E q o r q u k v k q p , ,

K p i t g f k g p v u	I o u " l " N k v t g
Proteose peptone	10.000
HM peptone B #	1.000
Sodium chloride	75.000
D-Mannitol	10.000
Phenol red	0.025
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Equivalent to Beef extract

F k t g e v k q p u

Suspend 96.0 grams in 1000 ml purified / distilled water. Heat if necessary to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C.

Note : This product contains 7.5% sodium chloride as one of its ingredients. On repeated exposure to air and absorption moisture sodium chloride has tendency to form lumps, therefore we strongly recommend storage in tightly closed containers in dry place away from bright light.

R t k p e k r i n g " C p f " K p v g t r t g v e v k q p

Mannitol Salt Broth is prepared as suggested by Chapman (1) and is used for the selective isolation of pathogenic Staphylococci. This medium is recommended for the detection and enumeration of coagulase-positive Staphylococci in milk (2) food (3) and other specimens. Mannitol Salt Broth is used for the isolation of presumptive pathogenic staphylococci. Pathogenic staphylococci ferment mannitol and produce a yellow coloured medium.

The medium contains beef extract and proteose peptone which makes it very nutritious as they provide essential growth factors and trace nutrients. Many other bacteria except Staphylococci are inhibited by 7.5% sodium chloride. Mannitol is the fermentable carbohydrate source. The differential action of the medium is attributed to D-Mannitol. *Staphylococcus aureus* ferments mannitol to produce yellow coloured medium. Most coagulase-negative species of Staphylococci and Micrococci do not ferment mannitol and therefore the medium remains red in colour. The colour of the medium is due to the reactivity of phenol red to the pH of the medium; phenol red is red at pH 8.4 and yellow at 6.8. Presumptive *Staphylococcus* showing yellow coloured medium should be further tested for production of coagulase.

A possible *S. aureus* must be confirmed by the coagulase test. Also the organism should be subcultured to a less inhibitory medium not containing excess salt to avoid the possible interference of salt with coagulase testing or other diagnostic tests (e.g. Nutrient Broth) (M002) (4). Few strains of *S. aureus* may exhibit delayed mannitol fermentation. Negative results should therefore be re-incubated for an additional 24 hours before being discarded (4).

V { r g " q h " u r g e k o g p

Clinical samples: pus, urine; Food and dairy samples, water samples.

Specimen Collection and Handling

) R U F O L Q L F D O V D P S O H V I R O O R Z D S S U R S U L D W H W H F K Q L T X H V I R U K D Q G O L Q
or food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines 12,1 .

At the same time, contaminated materials must be sterilized by autoclaving before discarding .

