



Moeller Decarboxylase Broth Base (Decarboxylase Broth Base, Moeller)

M393

Moeller Decarboxylase Broth Base with the addition of appropriate L-amino acid is used to differentiate bacteria on the basis of their ability to decarboxylate the amino acids.

Composition**

Ingredients	Gms / Litre
Peptone	5.000
Meat extract B#	5.000
Dextrose	0.500
Bromocresol purple	0.010
Cresol red	0.005
Pyridoxal	0.005
Final pH (at 25°C)	6.0±0.2

**Formula adjusted, standardized to suit performance parameters

#- Equivalent to Beef extract

Directions

Suspend 10.52 grams in 1000 ml distilled water. Add 10 gm. of L-Lysine, L-Arginine, L-Ornithine or other L-amino acids. When using DL-amino acids, use 2% concentration. Heat if necessary to dissolve the medium completely. When L-Ornithine is added, readjustment of the pH is required. Dispense in 5 ml amount in screw-capped tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 10 minutes.

Principle And Interpretation

Moeller Decarboxylase Broth Base is used for differentiating gram-negative enteric bacilli on the basis of their ability to decarboxylate amino acids. Moeller introduced the Decarboxylase Broth for detecting the production of lysine and ornithine decarboxylase and arginine dihydrolase (1). Prior to Moellers work, bacterial amino acid decarboxylases were studied by Gale (2) and Gale and Epps (3). Production of ornithine decarboxylase is a helpful criterion in differentiating *Klebsiella* and *Enterobacter* species. *Klebsiella* are nonmotile and do not produce ornithine decarboxylase while *Enterobacter* are motile and produce ornithine decarboxylase except *Enterobacter agglomerans* (4).

This medium contains meat extract B and peptone which provides nitrogenous and carbonaceous compounds, long chain amino acids and other essential nutrients for the growth of bacteria. Dextrose is the fermentable carbohydrate and pyridoxal is the co-factor for the decarboxylase enzyme. Bromo cresol purple and cresol red are the pH indicators in this medium. When the medium is inoculated with the dextrose fermenting bacteria, the pH is lowered due to acid production, which changes the colour of the indicator from purple to yellow. Acid produced stimulates decarboxylase enzyme. Decarboxylation of lysine yields cadaverine while putrescine is produced due to ornithine decarboxylation. Arginine is first hydrolyzed to ornithine which is then decarboxylated to form putrescine. Formation of these amines increases the pH of the medium, changing the colour of the indicator from yellow to purple. If the organisms do not produce the appropriate enzyme, the medium remains acidic, yellow in colour. Each isolate to be tested should also be inoculated into Moeller Decarboxylase Broth Base medium tube lacking the amino acid.

Inoculated tubes must be protected from air with a layer of sterile mineral oil. Exposure to air may cause alkalization at the surface of the medium which makes the test invalid.

Type of specimen

Clinical samples - Blood ; Food and dairy samples ; Water samples

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (4,5).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,2,8).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(3)

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

Limitations :

1. Some fastidious organisms may show delayed reaction.
2. Overlaying with mineral oil is essential for appropriate results.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to greenish yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Purple coloured, clear solution without any precipitate in tubes

Reaction

Reaction of 1.05% w/v aqueous solution at 25°C. pH : 6.0±0.2

pH

5.80-6.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for upto 4 days with addition of appropriate amino acids and overlaying with sterile mineral oil.

Cultural Response

Organism	Inoculum (CFU)	Arginine decarboxylation	Ornithine decarboxylation	Lysine decarboxylation
Cultural Response				
<i>Citrobacter freundii</i> ATCC 8090	50-100	variable reaction	variable reaction	negative reaction, yellow colour
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	negative reaction, yellow colour	positive reaction, purple colour	positive reaction, purple colour
<i>Escherichia coli</i> ATCC 25922	50-100	variable reaction	variable reaction	positive reaction, purple colour
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	negative reaction, yellow colour	negative reaction, yellow colour	positive reaction, purple colour
<i>Proteus mirabilis</i> ATCC 25933	50-100	negative reaction, yellow colour	positive reaction, purple colour	negative reaction, yellow colour
<i>Proteus vulgaris</i> ATCC 13315	50-100	negative reaction, yellow colour	negative reaction, yellow colour	negative reaction, yellow colour
<i>Salmonella Paratyphi A</i> ATCC 9150	50-100	delayed positive reaction/positive reaction, purple colour	positive reaction, purple colour	negative reaction, yellow colour

<i>Salmonella Typhi</i> ATCC 6539	50-100	delayed positive reaction / negative reaction	negative reaction, yellow colour	positive reaction, purple colour
<i>Serratia marcescens</i> ATCC 8100	50-100	negative reaction, yellow colour	positive reaction, purple colour	positive reaction, purple colour
<i>Shigella dysenteriae</i> ATCC 13313	50-100	negative reaction/ delayed positive reaction	negative reaction, yellow colour	negative reaction, yellow colour
<i>Shigella flexneri</i> ATCC 12022	50-100	negative reaction/ delayed positive reaction	negative reaction, yellow colour	negative reaction, yellow colour
<i>Shigella sonnei</i> ATCC 25931	50-100	variable reaction	positive reaction, purple colour	negative reaction, yellow colour

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Reference


1. Moeller V., 1955, Acta Pathol. Microbiol. Scand. 36:158.
 2. Gale G. F., 1940, Biochem. J., 34:392.
 3. Gale and Epps, 1943, Nature, 152:327.
 4. MacFaddin J., 1980, Biochemical Tests for Identification of Medical Bacteria, 2nd ed., Williams and Wilkins, Baltimore.
 5. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
 6. Downes F. P. and Ito K., (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.
 7. Greenberg A. E., Clesceri L. S. and Eaton A. D., (Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st ed., APHA, Washington, D.C.
 8. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
- . Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., A HA Inc., Washington, D.C.



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
B /4-6 , MIDC, Dindori, Nashik MH

www.himedialabs.com



CE Partner 4U ,Esdoornlaan 13, 3951

DB Maarn The Netherlands,

www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.