



## Toluidine Blue DNA Agar

M613I

Toluidine Blue DNA Agar is recommended for detection of thermostable deoxyribonuclease activity to establish speciation of *S.aureus* in contaminated foods. The composition and performance criteria of this medium are as per the specifications laid down in ISO 8870:2006(E) 83:2006(E).

### Composition\*\*

Ingredients	Gms / Litre
Deoxyribonucleic acid (DNA)	0.300
Calcium chloride	0.110
Sodium chloride	10.000
Toluidine blue	0.093
Tris (hydroxymethyl) amino methane	6.060
Agar	10.000
Final pH ( at 25°C)	9.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 26.56 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely and continue to boil for 1 to 2 minutes. Sterilization is not necessary. Dispense into sterile Petri plates.

### Principle And Interpretation

The growth of *Staphylococcus aureus* in foods represents a potential public health hazard since many strains of *S.aureus* produce enterotoxins that cause food poisoning if ingested. Numerous outbreaks of staphylococcal intoxication are associated with cheese, stimulating numerous studies on the incidence and behaviour of staphylococci in milk and cheese. . Toluidine Blue DNA Agar (M613) is recommended by APHA for detection of the thermostable deoxyribonuclease activity to establish the speciation of *S. aureus* in contaminated foods (1). Toluidine Blue DNA Agar (M613I) is also recommended by ISO Committee (2) with a slight modification in concentration of calcium chloride and toluidine blue.

DNA in the medium enables the detection of DNase activity by getting depolymerized and forming a clear zone around the microbial growth. Inclusion of toluidine blue aids in detection of DNase activity by the production of a visible bright rose-pink coloured reaction due to its metachromatic properties. Tris amino methane forms the buffering system. Sodium chloride and calcium chloride provide the ions and also maintains osmotic equilibrium.

### Quality Control

#### Appearance

Light yellow to light grey homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.0% Agar gel.

#### Colour and Clarity of prepared medium

Blue coloured clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 2.65% w/v aqueous solution at 25°C. pH : 9.0±0.2

#### pH

8.80-9.20

#### Cultural Response

M613I: 18 hours old BHI broth culture is heated in boiling water bath for 15 minutes and studied for thermonuclease activity. 5 mm cut well are cut in agar plates and is filled with 25-30µl of this culture and incubated at 35-37°C for 4 hrs (or it can also be incubated at 50°C for 2 hrs) and observed for results.

Organism	DNase activity
<i>Staphylococcus aureus</i>	positive
ATCC 12600	reaction, pink

halos extending  
1mm beyond  
the well  
*Staphylococcus epidermidis* negative  
*ATCC 14990* reaction

### Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

### Reference

1. Downes F. P. and Ito K., (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.
2. International Organization for Standardization ISO, 8870 :2006 (E), IDF, 83:2006 (E)

Revision : 02 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.