



## Phenol Red Dulcitol Broth

M617

Phenol Red Dulcitol Broth is used for dulcitol fermentation studies of microorganisms.

### Composition\*\*

Ingredients	Gms / Litre
Proteose peptone	10.000
Beef extract	1.000
Sodium chloride	5.000
Dulcitol	5.000
Phenol red	0.018
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 21.02 grams in 1000 ml distilled water, mix well. Heat if necessary to ensure complete dissolution. Distribute in fermentation tubes (tubes containing inverted Durham's tubes). Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

### Principle And Interpretation

Phenol Red Broth Medium is formulated as per Vera (2) and is recommended to determine the fermentation reaction of carbohydrates for the differentiation of microorganisms (3, 4, 5). Phenol Red Broth Medium with various carbohydrates serves as a differential medium by aiding in differentiation of various species and genera by their ability to ferment the specific carbohydrate, with the production of acid or acid and gas (6). Phenol Red Dulcitol Broth is used to study dulcitol fermentation in various bacteria.

Proteose peptone and beef extract serve as sources for carbon and nitrogen. Sodium chloride is the osmotic stabilizer. Phenol red is the pH indicator, which turns yellow at acidic pH i.e. on fermentation of dulcitol. Gas formation is seen in Durhams tubes. All of the *Enterobacteriaceae* grow well in this medium. In addition to producing a pH colour shift, the production of mixed acids, notably butyric acids, often results in a pungent, foul odour from the culture medium (1).

### Quality Control

#### Appearance

Light yellow to pink coloured homogeneous free flowing powder

#### Colour and Clarity of prepared medium

Red coloured clear solution without any precipitate

#### Reaction

Reaction of 2.1% w/v aqueous solution at 25°C. pH : 7.4±0.2

#### pH

7.20-7.60

#### Cultural Response

M617: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours .

Organism	Inoculum (CFU)	Growth	Acid	Gas
<b>Cultural Response</b>				
<i>Citrobacter freundii</i> ATCC 8090	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	Negative reaction, no colour change	Negative reaction

<i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
<i>Proteus vulgaris</i> ATCC 13315	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
<i>Salmonella Typhi</i> ATCC 6539	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	Positive reaction, yellow colour	Positive reaction
<i>Serratia marcescens</i> ATCC 8100	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
<i>Shigella flexneri</i> ATCC 12022	50-100	luxuriant	Negative reaction, no colour change	Negative reaction

### Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

### Reference

1. Koneman E. W., Allen S. D., Janda W.M., Schreckenberger P.C., Winn W.C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company
2. Vera H. D., 1950, Am. J. Public Health, 40, 1267
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification -Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
4. Finegold S. M. and Baron E. J., 1986, Bailey and Scotts Diagnostic Microbiology, 7th Ed., The C.V. Mosby Co., St. Louis.
5. Ewing W. H., 1986, Edwards and Ewings Identification of Enterobacteriaceae, 4th ed., Elsevier Science Publishing Co., Inc., New York.
6. MacFaddin J. F., 2000, Biochemical tests for Identification of Medical Bacteria, 3rd ed., Lippincott, Williams and Wilkins, Baltimore.

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.