



## Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Quality Control

### Appearance

Light yellow to light tan homogeneous free flowing powder

### Colour and Clarity of prepared medium

Light purple coloured, clear solution in tubes

### Reaction

Reaction of 1.6% w/v aqueous solution at 25°C. pH : 7.4±0.2

### pH

7.20-7.60

### Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 40-48 hours.

| Organism  | Inoculum (CFU)    | Growth         | Colour of medium            |
|---|-------------------|----------------|-----------------------------|
| <i>Escherichia coli</i> ATCC 25922 (00013*)           | 50-100            | good-luxuriant | purple                      |
| <i>Staphylococcus aureus</i> ATCC 25923 (00034*)      | 50-100            | none-poor      | light yellow                |
| <i>Staphylococcus epidermidis</i> ATCC 12228 (00036*) | 50-100            | fair-good      | purple/very slightly yellow |
| <i>Streptococcus pyogenes</i> ATCC 19615              | >=10 <sup>4</sup> | inhibited      |                             |

Key : \*Corresponding WDCM numbers.

## Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.