



Wilkins Chalgren Anaerobic Agar Base

M832

Wilkins Chalgren Anaerobic Agar Base is used for the selective isolation and cultivation of anaerobic bacteria and also for susceptibility testing of anaerobic bacteria by the agar dilution method.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Peptic digest of animal tissue	10.000
Yeast extract	5.000
Dextrose	1.000
Sodium chloride	5.000
L-Arginine	1.000
Sodium pyruvate	1.000
Hemin	0.005
Menadione	0.0005
Agar	10.000
Final pH (at 25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 43.0 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 50°C before adding antibiotics to be tested. Mix gently and pour into sterile Petri plates. For cultivation of anaerobes, aseptically add the rehydrated contents of 2 vials each of Non-Spore Anaerobic Supplement (FD001) or G. N. Spore Anaerobic Supplement (FD002) as desired to the sterile molten medium before pouring into sterile Petri plates.

Principle And Interpretation

Anaerobic bacteria are widespread in soil, marshes, lake and river sediments, oceans, sewage, food and animals. In humans, anaerobic bacteria normally are prevalent in the oral cavity around the teeth, in the gastrointestinal tract, in the orifices of the genitourinary tract and on the skin. Anaerobic infections in humans and various animals can involve virtually any organ under immunocompromised conditions (1). Also, anaerobic infections are often associated with tissue necrosis and abscess formation, leading to impaired delivery of antimicrobial agents in blood to the actual site of infection. This explains why anaerobic infections are often aggressively managed with debridement, aspiration and/or surgical removal of infected tissue. Because of the technical and interpretive difficulties associated with anaerobic susceptibility testing, presentation of definitive recommendations is difficult (2).

Wilkins Chalgren Anaerobic Agar Base, formulated by Wilkins and Chalgren (3), along with Brucella Agar Base is the preferred medium for agar dilution tests with anaerobes. This medium is also recommended for testing anaerobic bacteria (4, 5, 6). Wilkins Chalgren Agar need to be appropriately supplemented to support the growth of certain anaerobic bacteria. Hemin and Menadione (Vitamin K3) enhances the growth of *Bacteroides* species and *Prevotella melaninogenica*, respectively and many other species of gram-negative anaerobic rods (8, 6). The medium can also be supplemented with defibrinated or lysed blood for the growth of fastidious anaerobic bacteria (7).

Peptic digest of animal tissues and casein enzymic hydrolysate serve as sources of essential nutrients including carbon and nitrogen. Yeast extract provides vitamins and other growth factors like purines and pyrimidines that are essential for the growth of *P. melaninogenica*. Arginine serves as an amino acid source while pyruvate serves as an energy source. The medium can be made selective for non-sporing anaerobic bacteria and gram-negative anaerobic bacteria by addition of Non-Spore Anaerobic Supplement (FD001) and G. N. Spore Anaerobic Supplement (FD002) respectively.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel.

Colour and Clarity of prepared medium

Medium amber coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.3% w/v aqueous solution at 25°C. pH : 7.1±0.2

pH

6.90-7.30

Cultural Response

Cultural characteristics observed with added Non-spore Anaerobic Supplement (FD001) or G.N.Spore Anaerobic Supplement(FD002) under anaerobic condition, after an incubation at 35-37°C of 48 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response			
<i>Bacteroides fragilis</i> ATCC 25285	50-100	luxuriant	≥50%
<i>Clostridium perfringens</i> ATCC 12924	50-100	luxuriant	≥50%
<i>Escherichia coli</i> ATCC 25922	≥10 ³	inhibited	0%
<i>Prevotella melaninogenicus</i> ATCC 15930	50-100	luxuriant	≥50%

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

- Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company.
- Murray P. R., Baron E. J., Jorgensen J. H., Tenover F. C., Tenover F. C., (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.
- Wilkins T. D. and Chalgren S., 1976, Antimicrob. Agents Chemother., 10 : 926.
- King A., Phillips I., 1988, J. Antimicrob. Chemother., 21:425-438.
- Clinical and Laboratory Standards Institute, 2006, Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria, Approved standard M11-A3, CLSI, Villanova, Pa.
- Collee J. G., Fraser A. G., Marmion B. P., Simmons A., (Eds.), Mackie and McCartney Practical Medical Microbiology, 1996, 14th Edition, Churchill Livingstone.
- Gibbons R. J. and MacDonald J. B., 1960, J. Bacteriol., 80:164.
- Isenberg (Ed.), 2004, Clinical Microbiology Procedures Handbook, Vol.3, American Society for Microbiology, Washington. D.C.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.