



Blood Agar Base No. 2 with 1.2% Agar

M834A

Intended use

Recommended especially to permit the maximum recovery of fastidious pathogenic microorganisms without interfering with their haemolytic reactions.

Composition**

Ingredients	Gms / Litre
Proteose peptone	15.000
HL extract #	2.500
Yeast extract	5.000
Sodium chloride	5.000
Agar	12.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

-Equivalent to Liver extract

Directions

Suspend 19.75 grams in 500 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45 - 50°C and aseptically add 7% v/v sterile defibrinated blood.

For *Brucella* species: Add rehydrated contents of 1 vial of Brucella Selective Supplement (FD005) to 500 ml sterile molten base.

For *Campylobacter* species : Add rehydrated contents of 1 vial of Campylobacter Supplement - I (FD006) or Campylobacter Supplement - II (FD007) or Campylobacter Supplement - III (FD008) or Campylobacter Growth Supplement (FD009) to 500 ml sterile molten base.

For *Streptococcus* species: Add rehydrated contents of 1 vial of Strepto Supplement (FD031) to 500 ml sterile molten base. Mix well and pour into sterile Petri plates.

Principle And Interpretation

A fastidious organism is one with complete nutritional requirements, needing additional cellular building-block molecules in order to survive (4). This media is a highly nutritive, microorganisms producing haemolysin give visible haemolytic zones on this medium. It also serves as a differential medium for *Brucella* and *Campylobacter* species by adding different antibiotic supplements for the respective bacteria (1, 6). *Brucella* cultures are highly infective and must be handled with care. Incubate preferably in 5-10% carbon dioxide atmosphere. Comparative studies of horse, rabbit and sheep blood showed that sheep blood gave the clearest and most reliable colony and haemolysis characteristics at both 24 and 48 hours of incubation (7).

It can be used to prepare Chocolate Agar for the isolation of *Haemophilus* and *Neisseria* species. It can also be used for primary isolation of *Haemophilus* species, where horse blood is used for enrichment. Better results are obtained by spreading half of the horse blood agar plate with 2 drops of 10% saponin (9).

HL extract and yeast extract helps enhance the growth and haemolytic reactions of fastidious organisms like Streptococci and Pneumococci. Proteose peptone serves as the nitrogen source while HL extract and yeast extract provide essential carbon, vitamin, nitrogen and amino acid sources. Sodium chloride maintains the osmotic equilibrium. Supplementation with blood (5-10%) provides additional growth factors and also serves as basis for determining haemolytic reactions. Haemolytic patterns may vary with the source of animal blood or type of base medium used (3).

Type of specimen

Clinical material : blood and other pathological material; Food samples

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (2,3).

For food samples, follow appropriate techniques for sample collection and processing as per guidelines (5,8).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

- 1.Addition of sheep blood is recommended to detect haemolysis. This medium does not support the growth of *H.haemolyticus*
- 2.Addition of Horse blood or rabbit blood to base medium supports growth of *H.haemolyticus* but resemble beta-haemolytic Streptococci and hence must be confirmed.
3. Haemolytic pattern varies with the source of blood used.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.2% Agar gel

Colour and Clarity of prepared medium

Basal medium : Light amber coloured clear to slightly opalescent gel. After addition of 5% v/v sterile defibrinated blood : Cherry red coloured opaque gel forms in Petri plates.

Reaction

Reaction of 4.0% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20 - 7.60

Cultural Response

Cultural characteristics observed with added 5% w/v sterile defibrinated blood,after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth w/o blood	Recovery w/o blood	Growth with blood	Recovery with blood	Haemolysis
<i>Neisseria meningitidis</i> ATCC 13090	50-100	fair	40-50%	luxuriant	≥70%	none
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	good	50-70%	luxuriant	≥70%	beta
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	fair-good	40-50%	luxuriant	≥70%	alpha
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	fair-good	40-50%	luxuriant	≥70%	beta

Key : (*)Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10- 30°C in a tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

Reference

1. Hunter D. and Kearns M., 1977, Brit. Vet. J., 133:486.
2. Isenberg, H.D. Clinical Microbiology Procedures Handbook, 2nd Edition
3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual Clinical Microbiology, 11th Edition. Vol. 1.
4. Norton C. F., 1986, Microbiology, 2nd Edition, Addison-Wesley Publishing Company.
5. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
6. Skirrow M. B., 1977, B.M.J., ii: 9.
7. Snavey and Brahier, 1960, Am. J. Clin. Pathol., 33:511.
8. U.S. Food and Drug Administration, 1995, Bacteriological Analytical Manual, 8th Ed., AOAC International, Gaithersburg, Md.
9. Waterworth and Pamela M., 1955, Brit. J. Exp. Pathol., 36:186.


Revision : 04 / 2019



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg, Mumbai-86, MS, India



CE Partner 4U , Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.