



Technical Data

MacConkey HiCynth™ Agar w/o CV, NaCl and Sodium Taurocholate

MCD082

Recommended for cultivation and differentiation of enteric bacteria, restricting swarming of *Proteus* species from specimens such as urine which may contain large number of *Proteus* species as well as potentially pathogenic Gram-positive organisms.

Composition**

Ingredients	Gms / Litre
HiCynth™ Peptone No.3*	23.000
Lactose	10.000
Synthetic detergent	2.000
Neutral red	0.040
Agar	20.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

*Chemically defined peptone

Directions

Suspend 55.04 grams of medium in 1000 ml distilled water. Heat to boiling with gentle swirling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Avoid overheating. Cool to 45-50°C. Mix well and pour into sterile Petri plates. The surface of the medium should be dry when inoculated.

Principle And Interpretation

MacConkey Agar is the earliest selective and differential medium for cultivation of enteric microorganisms from a variety of clinical specimens (1, 2). Subsequently MacConkey Agar and Broth have been recommended for use in microbiological examination of foodstuffs (3) and for direct plating / inoculation of water samples for coliform counts (4). These media are also accepted by the Standard Methods for the Examination of Milk and Dairy Products (5) and pharmaceutical preparations (6). MacConkey HiCynth™ Agar w/o CV, NaCl, and Sodium Taurocholate is prepared by completely replacing animal or vegetable peptones with chemically defined peptones to avoid BSE/TSE risks associated with animal peptones.

Gram-negative bacteria usually grow well on the medium and are differentiated by their ability to ferment lactose. Lactose fermenting strains grow as red or pink. The red colour is due to production of acid from lactose, absorption of neutral red and a subsequent colour change of the dye when the pH of medium falls below 6.8. Lactose non-fermenting strains, such as *Shigella* and *Salmonella* are colourless and transparent and typically do not alter appearance of the medium. *Yersinia enterocolitica* may appear as small, non-lactose fermenting colonies after incubation at room temperature.

The medium contains HiCynth™ Peptone No.3 which provides nitrogenous and carbonaceous compounds, long chain amino acids, vitamins and other essential nutrients. The selective action of this medium is attributed to synthetic detergent, which are inhibitory to most species of gram-positive bacteria. It is a modification of the original formulation with the exclusion of crystal violet like the regular MacConkey Agar.

Quality Control

Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 2.0% Agar gel.

Colour and Clarity of prepared medium

Orange red coloured clear to slightly opalescent gel forms in Petri plates

Please refer disclaimer Overleaf.

Reaction

Reaction of 5.5% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20-7.60

Cultural Response

Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony
Cultural Response <i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	≥50%	pink to red
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	≥50%	pale pink to red
<i>Enterococcus faecalis</i> ATCC 29212	50-100	fair to good	30-40%	pale pink to red
<i>Proteus vulgaris</i> ATCC 13315	50-100	luxuriant	≥50%	colourless
<i>Salmonella Paratyphi A</i> ATCC 9150	50-100	luxuriant	≥50%	colourless
<i>Shigella flexneri</i> ATCC 12022	50-100	fair to good	30-40%	colourless
<i>Salmonella Paratyphi B</i> ATCC 8759	50-100	luxuriant	≥50%	colourless
<i>Salmonella Enteritidis</i> ATCC 13076	50-100	luxuriant	≥50%	colourless
<i>Salmonella Typhi</i> ATCC 6539	50-100	luxuriant	≥50%	colourless
<i>Staphylococcus aureus</i> ATCC 25923	50-100	fair-good	30-40%	pale pink -red

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. MacConkey, 1900, The Lancet, ii:20.
2. MacConkey, 1905, J. Hyg., 5:333.
3. Downes F. P and Ito K. (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th ed., APHA, Washington, D.C.
4. Greenberg A. E., Clesceri L. S. and Eaton A. D., (Eds.), 2012, Standard Methods for the Examination of Water and Wastewater, 22nd ed., APHA, Washington, D.C.
5. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.
6. The United States Pharmacopoeia XXI and the National Formulary, 16th ed., 1985, United States Pharmacopoeial Convention, Inc., Washington, D.C.

Revision : 00 / 2015

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.