

MUG Plate Count HiCynth™ Agar

MCD1194

MUG Plate Count HiCynth™ Agar is used for determination of plate count of microorganisms in milk and other dairy products by fluorogenic method.

Composition**

Ingredients	Gms / Litre
HiCynth™ Peptone No.3*	5.000
HiCynth™ Peptone No.5*	2.500
Dextrose	1.000
4-Methylumbelliferyl β-D-Glucuronide (MUG)	0.100
Agar	15.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

*Chemically defined peptones

Directions

Suspend 23.6 grams in 1000 ml distilled water. Heat gently to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Plate Count Agar is a general-purpose cultivation medium used for a wide variety of organisms and is recommended by APHA (1, 2, 3) and AOAC (4).

MUG Plate Count HiCynth™ Agar, which is Plate Count HiCynth™ Agar supplemented with MUG, is used for determining plate count of microorganisms in milk and other dairy products by fluorogenic method. It is prepared by completely replacing animal or vegetable peptones with chemically defined peptones to avoid BSE/TSE risks associated with animal peptones. The medium does not contain any inhibitor or pH indicator. It is used to determine the total microbial count of milk, dairy products (1), water (2) and other materials. Organism like *Escherichia coli* can be identified by the formation of fluorescent colonies visualized on exposure to UV light (366nm).

HiCynth™ Peptone No.3 and HiCynth™ Peptone No.5 provide nitrogenous and carbonaceous compounds, long chain amino acids and vitamin B complex.

Dextrose serves as energy source. MUG is cleaved by the enzyme beta-glucuronidase to release 4-methylumbelliferone, which produces a visible fluorescence under long wave UV light.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 2.36% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Fluorescence (under UV)
Cultural Response				
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	≥70%	positive
<i>Staphylococcus aureus</i> ATCC 25923	50-100	luxuriant	≥70%	negative
<i>Bacillus subtilis</i> ATCC 6633	50-100	luxuriant	≥70%	negative
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	luxuriant	≥70%	negative
<i>Enterococcus faecalis</i> ATCC 29212	50-100	luxuriant	≥70%	negative
<i>Lactobacillus casei</i> ATCC 9595	50-100	luxuriant	≥70%	negative

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

1. Richardson G., (Ed.), 1985, Standard Methods for the Examination of Dairy Products, 15th Ed., APHA, Washington, D.C.
2. Greenberg A. E., Trussell R. R. and Clesceri L. S., (Eds.), 1985, Standard Methods for the Examination of Water and Wastewater, 16th Ed., APHA, Washington, D.C.
3. Downes F.P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.
4. Horwitz, (Ed.), 2000, Official Methods of Analysis of AOAC International, 17th Ed. Vol. I, AOAC International, Gaithersburg, Md.

Revision : 00/ 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.