

Pseudomonas HiCynth™ Agar (For Fluorescein)

MCD120

Pseudomonas HiCynth™ Agar (For Fluorescein) is recommended for the detection of fluorescein production by *Pseudomonas* species.

Composition**

Ingredients	Gms / Litre
HiCynth™ Peptone No.3*	15.000
HiCynth™ Peptone No.5*	5.000
Dipotassium phosphate	1.500
Magnesium sulphate	1.500
Agar	15.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

*Chemically defined peptones.

Directions

Suspend 38 grams in 1000 ml distilled water containing 10 ml glycerol. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C . Mix well and pour into sterile Petri plates.

Principle And Interpretation

Pseudomonas Agar (For Fluorescein) is based on the formula described by King et al (1) and as modified in the U.S. Pharmacopeia (2) for the detection of fluorescein production a water soluble, chloroform insoluble fluorescent pigment by *Pseudomonas* species (3). Pseudomonas HiCynth™ Agar (For Fluorescein) is a modification of Pseudomonas Agar (For Fluorescein) wherein animal based peptone is replaced with chemically defined peptone to avoid BSE/TSE risks associated with animal peptone. The medium enhances the elaboration of fluorescein by *Pseudomonas* and inhibits the pyocyanin formation. The fluorescein pigment diffuses from the colonies of *Pseudomonas* into the agar and shows yellow fluorescent colouration. Some *Pseudomonas* strains produce small amounts of pyocyanin resulting in a yellow-green colouration.

HiCynth™ Peptone No.3 and HiCynth™ Peptone No.5 provide the essential nitrogenous and carbonaceous nutrients, carbon, amino acids, vitamins and trace elements for the growth of *Pseudomonas*.

Dipotassium phosphate buffers the medium while magnesium sulphate provides necessary cations for the activation of fluorescein production. Salt concentration exceeding 2% affects pigment production. UV illumination may be bactericidal, so make sure that there is good growth before placing culture under UV light (3).

A pyocyanin-producing *Pseudomonas* strain will usually also produce fluorescein. It must, therefore, be differentiated from other simple fluorescent pseudomonads by other means. Temperature can be a determining factor as most other fluorescent strains will not grow at 35°C. Rather, they grow at 25-30°C (2).

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.8% w/v aqueous solution (containing 1% v/v glycerol) at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural characteristics observed with added 1% glycerol after an incubation at 35-37°C for 18-24 hours.

Cultural Response

Please refer disclaimer Overleaf.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
Cultural Response				
<i>Pseudomonas aeruginosa</i> ATCC 17934	50-100	luxuriant	>=70%	greenish yellow
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	luxuriant	>=70%	greenish yellow
<i>Pseudomonas aeruginosa</i> ATCC 9027	50-100	luxuriant	>=70%	greenish yellow

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. King, Ward and Raney, 1954, J. Lab. Clin. Med., 44 : 301.
2. MacFaddin J., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
3. The United States Pharmacopoeia, 2008, USP31/NF26, The United States Pharmacopoeial Convention, Rockville, MD.

Revision : 00 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.