

## Bile Esculin Azide HiCynth™ Agar

MCD493

Bile Esculin Azide HiCynth™ Agar is a selective medium used for isolation and presumptive identification of faecal Streptococci.

### Composition\*\*

Ingredients	Gms / Litre
HiCynth™ Peptone No.3*	25.000
HiCynth™ Peptone No.5*	5.000
Synthetic detergent No.II	5.000
Esculin	1.000
Ferric ammonium citrate	0.500
Sodium chloride	5.000
Sodium azide	0.150
Agar	15.000
Final pH ( at 25°C)	7.1±0.2

\*\*Formula adjusted, standardized to suit performance parameters

\*Chemically defined peptones

### Directions

Suspend 56.65 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

*Caution: Sodium azide has a tendency to form explosive metal azides with plumbing materials. It is advisable to use enough water to flush off the disposables.*

### Principle And Interpretation

Group D Streptococci possess the group D lipoteichoic acid antigen in their cell walls. Former Group D species, which are predominant normal inhabitants of the human gastrointestinal tract, are termed as faecal Streptococci or Enterococci (1). The unique ability of Enterococci to split esculin was reported by Meyer and Schonfeld (2). Enterococci and Group D Streptococci hydrolyse esculin to esculin and dextrose, which reacts with ferric citrate producing brownish black precipitate (3). The use of esculin hydrolysis in identification of Enterococci was first cited by Rochaix (4). Bile Esculin Agar was originally formulated by Swan (6) which is modified to synthetic media, for the isolation and identification of Group D Streptococci from food. Facklam and Moody (7, 8) further reported that using Bile Esculin Agar, Group D Streptococci could be differentiated from non Group D Streptococci. Bile Esculin Agar was also shown to aid differentiation of *Enterobacteriaceae*, *Klebsiella*, *Enterobacter*, *Serratia* from other *Enterobacteriaceae* genera (9) on the basis of esculin hydrolysis. However, other tests such as salt tolerance should be performed for identifying Enterococci (5).

Bile Esculin Azide HiCynth™ Agar is a modification of Bile Esculin Agar wherein animal or vegetable based peptones are replaced with chemically defined peptones to avoid BSE/TSE risks associated with animal peptones. (6, 8) as per Isenberg (10). In this medium the bile concentration is reduced and additional sodium azide is incorporated.

HiCynth™ Peptone No.3 and HiCynth™ Peptone No.5 serves as source of carbon, nitrogen, amino acids, vitamins and essential growth nutrients. Synthetic detergent II and Sodium azide inhibits most of the other accompanying bacteria. Esculin in the medium is hydrolyzed to esculin and dextrose. Esculetin reacts with ferric citrate to form a dark brown or black complex, visualized as a zone of black precipitate around the colonies. If the media is dispensed in tubes in the form of slants, a positive reaction is indicated by blackening of more than half of the slant within 24-48 hours. If blackening is totally absent or if less than half of the slant is blackened within 24-48 hours, the test is negative. Viridans Streptococci sometimes exhibit a weak positive reaction. Also, *Leuconostoc*, *Pediococcus*, *Lactococcus* species causing human infections give a positive bile esculin test (11). To enhance the growth of Enterococci, Bile Esculin HiCynth™ Agar can be supplemented with 50ml/l horse serum (3).

Suspected water samples are filtered using membrane filters. These membrane filters are aseptically placed on Slanetz and Bartely HiCynth™ Medium. Red or maroon coloured colonies observed after incubation are further confirmed by aseptically transferring the membrane filter on to Bile Esculin Azide HiCynth™ Agar plate preheated to 44°C. Incubation at 44 ± 0.5°C for 2 hours is done following the inoculation. All typical colonies exhibiting a brown black colouration in the surrounding medium are counted as intestinal Enterococci (11).

Alternatively Bile Esculin Azide HiCynth™ Agar can also be used for direct isolation of Enterococci (without membrane filter), by incubation at 35-37°C for 18-24 hours.

## Quality Control

### Appearance

Cream to yellow homogeneous free flowing powder

### Gelling

Firm, comparable with 1.5% Agar gel

### Colour and Clarity of prepared medium

Amber coloured, clear to slightly opalescent gel with a bluish tinge forms in Petri plates.

### Reaction

Reaction of 5.67% w/v aqueous solution at 25°C. pH : 7.1±0.2

### pH

6.90-7.30

### Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

### Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Esculin Hydrolysis
<b>Cultural Response</b> <i>Enterococcus faecalis</i> ATCC 50-100 29212		luxuriant	≥50%	positive reaction, blackening of medium around the colony
<i>Escherichia coli</i> ATCC 25922	≥10 <sup>3</sup>	inhibited	0%	
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good	40-50%	negative reaction
<i>Proteus mirabilis</i> ATCC 25933	50-100	good	40-50%	negative reaction
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	none-poor	≤10%	negative reaction

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

## Reference

- Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4 th Ed., J. B. Lippincott Company
- Meyer and Schonfeld, 1926, Zentralbl. Bakteriol, Parasitenk. Infektionskr. Hyg. Abt. Orig. 99:402.
- MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
- Rochaix, 1924, Comt. Rend. Soc. Biol., 90:771.
- Facklam R., 1973, Appl. Microbiol., 26:138.
- Swan, 1954, J. Clin. Pathol., 7:160.
- Facklam R., 1972, Appl. Microbiol., 23:1131.
- Facklam R. R and Moody M. D., 1970, Appl. Microbiol., 20(2):245.
- Edberg S. C., Pittman S., and Singer J. M., 1977, J. Clin. Microbiol., 6:111.
- Isenberg, 1970, Clin. Lab. Forum, July.
- Murray P. R., Baron E. J., Jorgensen J. H., Tenover F. C., Tenover F. C., (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.