



Medium 10. Levin Eosin- Methylene Blue Agar

MM022

Levin Eosin Methylene Blue Agar Medium is recommended for the isolation, enumeration and differentiation of members of *Enterobacteriaceae* in accordance with Indian Pharmacopoeia..

Composition**

Ingredients	Gms / Litre
Pancreatic digest of gelatin	10.000
Dibasic potassium phosphate	2.000
Lactose	10.000
Eosin - Y	0.400
Methylene blue	0.065
Agar	15.000
pH after sterilization (at 25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 37.46 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes or as per validated cycle. AVOID OVERHEATING. Cool to 50°C and shake the medium in order to oxidize the methylene blue (i.e. restore its blue colour) and to suspend the precipitate, which is an essential part of the medium. Mix well before pouring into sterile Petri plates.

Precaution : Store the medium away from light to avoid photooxidation.

Principle And Interpretation

Levin Eosin Methylene Blue Agar was developed by Levine (1,2) and is used for the differentiation of *Escherichia coli* and *Enterobacter aerogenes* and also for the rapid identification of *Candida albicans* . This medium is recommended for the detection, enumeration and differentiation of members of the coliform group by American Public Health Association and Indian Pharmacopoeia (3, 4, 5, 6).

Eosin-Y and methylene blue make the medium slightly selective and inhibit certain gram-positive bacteria. Both dyes act as indicator and inhibiting agent. These dyes differentiate between lactose fermenters and non-fermenters. Eosin Y and methylene blue forms a complex at acidic pH which acts as inhibiting agent. Some gram-positive bacteria such as faecal Streptococci, yeasts grow on this medium and form pinpoint colonies. Essential nutrients and growth factors is provided by pancreatic digest of gelatin. Phosphates act as good buffering agent. *E.coli* forms colonies with green metallic sheen, indicating strong lactose fermentation. Weld (7,8) proposed the use of Levine EMB Agar, with added Chlortetracycline hydrochloride, for the rapid identification of *Candida albicans* in clinical specimens. A

positive identification of *Candida albicans* can be made after 24 48 hours incubation at 35 - 37°C in 10% carbon dioxide atmosphere, from specimens such as faeces, oral and vaginal secretions and nail or skin scraping etc. However, the typical appearance is variable.

Quality Control

Appearance

Light pink to purple homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Reddish purple with greenish cast coloured opalescent gel with finely dispersed precipitate forms in Petri plates.

pH

6.90-7.30

Cultural Response

Growth Promotion is carried out in accordance with IP. Cultural response was observed after an incubation at 36-38°C for 18-24 hours. Recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar.

Cultural Response

Organism	Inoculum (CFU)	Observed Lot value (CFU)	Recovery	Colour of Colony	Incubation temperature	Incubation period
Test for specified microorganism						
<i>Escherichia coli</i> ATCC 8739	50 -100	25 -100	>=50 %	blue-black colonies with metallic sheen	36 -38 °C	18 -24 hrs
<i>Escherichia coli</i> NCTC 9002	50 -100	25 -100	>=50 %	blue-black colonies with metallic sheen	36 -38 °C	18 -24 hrs

Additional Microbiological testing

<i>Escherichia coli</i> ATCC 25922	50 -100	25 -100	>=50 %	blue-black colonies with metallic sheen	36 -38 °C	18 -24 hrs
<i>Enterobacter aerogenes</i> ATCC 13048	50 -100	25 -100	>=50 %	pink to red	36 -38 °C	18 -24 hrs
<i>Salmonella Typhimurium</i> ATCC 14028	50 -100	25 -100	>=50 %	colourless	36 -38 °C	18 -24 hrs
<i>Pseudomonas aeruginosa</i> ATCC 9027	50 -100	25 -100	>=50 %	colourless	36 -38 °C	18 -24 hrs
<i>Enterococcus faecalis</i> ATCC 29212	>=10 ³	0	0%		36 -38 °C	18 -72 hrs
<i>Staphylococcus aureus</i> ATCC 6538	>=10 ³	0	0%		36 -38 °C	18 -72 hrs
<i>Candida albicans</i> ATCC 10231	50 -100	25 -100	>=50 %	colourless	36 -38 °C	18 -24 hrs
<i>Saccharomyces cerevisiae</i> ATCC 9763	50 -100	0 -10	0 -10 %	cream	36 -38 °C	18 -24 hrs

Storage and Shelf Life

Store below 30°C in tightly closed container and prepared medium at 2 - 8°C and away from light. Use before expiry date on the label.

Reference

1. Levine M., 1918, J. Infect. Dis., 23:43.
2. Levine M., 1921, Bull. 62, Iowa State College Engr. Exp. Station.
3. Eaton A. D., Clesceri L. S. and Greenberg A W.,(Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st ed., APHA, Washington, D.C.
4. Wehr H M and Frank J H., 2004, Standard Methods for the Examination of Dairy Products, 17th ed., APHA Inc., Washington, D.C.
5. Downes F P and Ito K. (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th ed., APHA, Washington, D.C.
6. Indian Pharmacopoeia, 2007, Govt. of India, Ministry of Health and Family Welfare, New Delhi.
7. Weld J. T., 1952, Arch. Dermat. Syph., 66:691.
8. Weld J. T., 1953, Arch. Dermat. Syph., 67(5):433.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.