



Baird-Parker Agar Medium

MU043

Baird-Parker Agar Medium is recommended for the selective isolation and enumeration of coagulase positive Staphylococci from food and other materials in accordance with United States Pharmacopoeia.

Composition**

Ingredients	Gms / Litre
Pancreatic digest of casein	10.000
Beef extract	5.000
Yeast extract	1.000
Glycine	12.000
Sodium pyruvate	10.000
Lithium chloride	5.000
Agar	20.000
pH after sterilization (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 63 grams in 950 ml purified /distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and add aseptically 50 ml concentrated Egg Yolk Emulsion (FD045) and 10 ml sterile 1% Potassium Tellurite solution (FD052). Mix well before pouring into sterile Petri plates..

Warning : Lithium Chloride is harmful. Avoid all bodily contact and inhalation of vapours. On contact with skin wash with plenty of water immediately.

Principle And Interpretation

This medium was first described in 1952. This medium was developed by Baird-Parker (1,2) from the Tellurite-Glycine formulation of Zebovitz et al (3) for selective isolation of *Staphylococcus aureus* from foods. *Staphylococcus* species are common contaminants in food, dairy, pharmaceutical and cosmetics related products (9). This medium is recommended for microbial limit tests of non-sterile pharmaceutical products and to detect *S.aureus*. Baird Parker Agar Medium was reported to be the best medium for selective detection of coagulase positive and enterotoxigenic *Staphylococcus* (4). This medium was found to be less inhibitory to *S.aureus* than other media, at the same time being more selective (5, 6). Subsequently it was officially adapted by the AOAC and is also recommended in United States Pharmacopoeia for use in Microbial limit test (7, 8).

„Beef extract, yeast extract and pancreatic digest of casein provides essential minerals, vitamin and other growth requirements. Sodium pyruvate protects injured cells and helps recovery. Lithium chloride and potassium tellurite inhibit most of contaminating microflora except *S.aureus*. Glycine, pyruvate enhances growth of *Staphylococcus*. With the addition of egg yolk the medium becomes yellow and opaque.

„Proteolytic bacteria produce a clear zone around colony in egg yolk containing media also known as Lecithinase reaction. A clear zone and grey-black colonies on this medium are diagnostic for coagulase positive Staphylococci. Upon further incubation, an opaque zone is developed around colonies, which can be due to lipolytic activity. Identity of *Staphylococcus aureus* isolated on Baird-Parker Agar must be confirmed with a coagulase reaction.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 2.0% agar gel.

Colour and Clarity of prepared medium

Basal medium: Yellow coloured clear to slightly opalescent gel. After addition of Egg Yolk Emulsion and Tellurite Emulsion: Yellow coloured opaque gel forms in Petri plates.

Reaction

After sterilization, reaction of 6.3% w/v aqueous solution. pH : 6.8±0.2

pH

6.60-7.00

Cultural Response

Growth Promotion is carried out in accordance USP. Cultural response was observed after an incubation at 30-35°C for 24-48 hours. Recovery rate is considered as 100% for bacteria growth on Soybean Casein Digest Agar.

Cultural Response

Organism	Inoculum (CFU)	Growth	Observed Lot value (CFU)	Recovery	Colour of colony	Lecithinase
Growth Promoting						
<i>Staphylococcus aureus</i> ATCC 6538	50 -100	luxuriant	25 -100	≥50 %	grey-black shiny	Positive, opaque zone around the colony
Additional Microbiological testing						
<i>Staphylococcus aureus</i> ATCC 25923	50 -100	luxuriant	25 -100	≥50 %	grey-black shiny	Positive, opaque zone around the colony
<i>Proteus mirabilis</i> ATCC 25933	50 -100	good - luxuriant	25 -100	≥50 %	brown - black	Negative
<i>Micrococcus luteus</i> ATCC 10240	50 -100	poor - good	15 -40	30 -40 %	shades of brown-black (very small)	Negative
<i>Staphylococcus epidermidis</i> ATCC 12228	50 -100	poor - good	15 -40	30 -40 %	black	Negative
<i>Bacillus subtilis</i> ATCC 6633	50 -100	none - poor	0 -10	0 -10 %	dark brown matt	Negative
<i>Escherichia coli</i> ATCC 8739	50 -100	none- poor	0 -10	0 -10 %	large brown black	Negative
<i>Escherichia coli</i> ATCC 25922	50 -100	none- poor	0 -10	0 -10 %	large brown black	Negative
<i>Escherichia coli</i> NCTC 9002	50 -100	none- poor	0 -10	0 -10 %	large brown black	Negative

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

1. Baird-Parker, A.C. 1962, J. Appl. Bact., 25: 12.
2. Baird-Parker, A.C. and Davenport, E., 1965, J. Appl. Bact., 28: 390.
3. Zebowitz, E., Evans J.B. & Niven C.F., (1955), J. Bact; 70:686.
4. Niskanean A and Aalto M, App. Env. Microbiol., 1978, 35:1233.
5. Tardio and Baer, 1971, J. Assoc. Off. Anal. Chem., 54:728.
6. Baer, 1971, J. Assoc. Off. Anal. Chem., 54:732.
7. The United States Pharmacopoeia, 2009, US Pharmacopoeial Convention Inc. Twinbrook Parkway, Rockville, M.D..
8. J. Assoc. off. Anal. Chem, 1971, 54:401.
9. Bacteriological Analytical Manual, 8th Edition, Revision A, 1998. AOAC, Washington D.C.

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.