

# Murashige and Skoog Medium

(Finer & Nagasawa Modification)  
With Macroelements and Microelements  
Without Vitamins, Sucrose and Agar

**Product Code: PT051**

## Product Description :

Murashige and Skoog Medium (MS) was originally formulated by Murashige and Skoog in 1962 to optimize tobacco callus bioassay system for facilitating the study of cytokinins. Since then, it is widely used for micro propagation, organ culture, callus culture and suspension culture.

Murashige and Skoog Medium is a nutrient blend of inorganic salts that provides all essential macroelements and microelements. Increased potassium nitrate serves as a source of nitrogen which aids in growth of callus significantly, Potassium dihydrogen phosphate serves as a source of phosphate. Microelements like Boron, Manganese, Molybdenum, Copper, Iron and Zinc enhance the catabolism in plants. Boron plays a key role in carbohydrate metabolism.

The product is plant tissue culture tested but it is the sole responsibility of the user to ensure the suitability of the medium for individual species.

## Composition :

Ingredients	mg/L
<b>MACROELEMENTS</b>	
Ammonium nitrate	825.000
Calcium chloride	332.200
Magnesium sulphate	180.690
Potassium nitrate	3040.000
Potassium phosphate monobasic	170.000
<b>MICROELEMENTS</b>	
Boric acid	6.200
Cobalt chloride hexahydrate	0.025
Copper sulphate pentahydrate	0.025
EDTA disodium salt dihydrate	37.300
Ferrous sulphate heptahydrate	27.800
Manganese sulphate monohydrate	16.900
Molybdic acid (sodium salt)	0.213
Potassium Iodide	0.830
Zinc sulphate heptahydrate	8.600
<b>Total(gms/litre)</b>	<b>4.6</b>

## Material required but not provided :

- Autoclaved distilled water
- MS Vitamins (VP021/ PL022)
- Sucrose (PCT0607)
- Plant growth regulators
- Gelling agents like Agar (PCT0901) or CleriGel™ (PCT0903)
- 1N NaOH/HCl

## Precautions :

- Ensure appropriate pH of the medium before addition of gelling agent as acidic pH will lead to decreased gelation resulting in semi solid flowing gel while alkaline pH will lead to formation of hardened gel.
- Use of Distilled water / Tissue culture grade water is recommended for media preparation as tap water or lower grade water may lead to salt precipitation and improper gelation.
- Avoid preparation of concentrated solutions, as it will lead to precipitation of salts.

## Directions :

- Reconstitute medium by adding required quantity of powder in two-third of total volume with constant, gentle stirring till the medium gets completely dissolved.
- Add heat stable supplements prior to autoclaving.
- Make up the final volume with distilled water.
- Adjust the pH of the medium to  $5.75 \pm 0.5$  using 1N NaOH/HCl.
- Add gelling agent and heat the medium to boiling till complete dissolution of gelling agent.
- Sterilize the medium by autoclaving at 15 lbs and 121°C for 15 min.
- Cool the autoclaved medium to about 45°C before adding heat labile supplements.
- Aseptically dispense the desired amount of medium under a laminar airflow unit in sterile culture vessels.

## Quality Control:

### Appearance

White to off-white, homogenous, free flowing powder

### Solubility

4.6 gms/litre soluble in distilled water

### Colour and Clarity

Colourless to light yellow, clear solution

### pH at 25°C

3.50 - 4.50

### Plant Tissue Culture Test

The growth promoting properties of medium is assessed by providing plant cultures with relative humidity of about  $60\% \pm 2\%$ , temperature  $22^{\circ}\text{C} \pm 2^{\circ}\text{C}$  and photoperiod of about 16:8. The plant species showed actively growing callus and shoots with no structural, necrotic and toxic deformity.

## Storage and Shelf Life:

- The plant tissue culture medium powder is extremely hygroscopic and must be stored at  $2-8^{\circ}\text{C}$  in air tight containers.
- Preferably, entire content of each package should be used immediately after opening.
- Use before the expiry date.

Revision : 01 / 2017

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic , research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.