

Vacin and Went Medium

With Thiamine, Sucrose and Agar

Product Code: PT082

Product Description :

Vacin and Went medium consists of the macroelements and microelements as described by Vacin and Went in 1949. The medium was developed for the *in vitro* culture of orchid species.

The formulation is a nutrient blend of inorganic salts, vitamin, carbohydrate and gelling agent. Potassium nitrate along with ammonium sulphate serve as sources of nitrogen and induces organogenesis. Calcium phosphate and potassium dihydrogen phosphate provide phosphate and enhance protocorm like bodies formation. Microelements like manganese and iron play a key role in the metabolism and enhance proliferation in the plant tissues. Thiamine hydrochloride acts as an important enzymatic cofactor in primary and secondary processes like glycolysis and TCA cycle.

The product is plant tissue culture tested but it is the sole responsibility of the user to ensure the suitability of the medium for individual species.

Composition :

Ingredients	mg/L
MACROELEMENTS	
Ammonium sulphate	500.000
Calcium phosphate	200.000
Magnesium sulphate	122.087
Potassium nitrate	525.000
Potassium phosphate monobasic	250.000
MICROELEMENTS	
Ferric tartarate	26.290
Manganese sulphate	5.683
VITAMINS	
Thiamine hydrochloride	0.400
CARBOHYDRATE	
Sucrose	20000.000
GELLING AGENT	
Agar	8000.000
Total(gms/litre)	29.6

Material required but not provided :

- Autoclaved distilled water
- Plant growth regulators
- 1N NaOH/HCl

Precautions :

- Ensure appropriate pH of the medium before addition of gelling agent as acidic pH will lead to decreased gelation resulting in semi solid flowing gel while alkaline pH will lead to formation of hardened gel.
- Use of Distilled water/Tissue culture grade water is recommended for media preparation as tap water or lower grade water may lead to salt precipitation and improper gelation.
- Avoid preparation of concentrated solutions, as it will lead to precipitation of salts.

Directions :

- Reconstitute medium by adding required quantity of powder in two-third of total volume with constant, gentle stirring till the medium gets completely dissolved.
- Add heat stable supplements prior to autoclaving.
- Make up the final volume with distilled water.
- Adjust the pH of the medium to 5.75 ± 0.5 using 1N NaOH/HCl.
- Heat the medium to boiling till complete dissolution of gelling agent.
- Sterilize the medium by autoclaving at 15 lbs and 121°C for 15 min.
- Cool the autoclaved medium to about 45°C before adding heat labile supplements.
- Aseptically dispense the desired amount of medium under a laminar airflow unit in sterile culture vessels.

Quality Control:

Appearance

White to off-white, homogenous, free flowing powder

Solubility

29.6 gms/litre soluble after boiling in distilled water

Colour and Clarity

Colourless to light yellow solution, hazy gel is formed on cooling

Gelling

Firm gel formed at pH: 5.75 ± 0.5

pH at 25°C

5.40 - 6.40

Plant Tissue Culture Test

The growth promoting properties of medium is assessed by providing plant cultures with relative humidity of about $60\% \pm 2\%$, temperature $22^\circ\text{C} \pm 2^\circ\text{C}$ and photoperiod of about 16:8. The plant species showed actively growing callus and shoots with no structural, necrotic and toxic deformity.

Storage and Shelf Life:

- The plant tissue culture medium powder is extremely hygroscopic and must be stored at $2-8^\circ\text{C}$ in air tight containers.
- Preferably, entire content of each package should be used immediately after opening.
- Use before the expiry date.

Revision : 01 / 2017

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic , research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.