



HiVeg™ Hydrolysate

RM014V

Principle And Interpretation

HiVeg™ Hydrolysate is prepared by enzymic hydrolysis of vegetable proteins to suit cultural response comparable with Tryptone, Type - I (Casitose, Type-1) (RM014). Recommended for the cultivation of a wide variety of organisms for indole production and fermentation studies. And also for bulk production of enzymes and antibiotics.

Quality Control

Appearance

Light yellow to yellow, may have a slight green tinge homogenous free flowing powder, having characteristic odour of protein, derived from vegetable source.

Solubility

Freely soluble in distilled/ purified water, insoluble in alcohol.

Clarity

1% w/v aqueous solution is clear without any haziness after autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Reaction

Reaction of 2% w/v aqueous solution at 25°C.

pH

5.50- 7.50

Microbial Load:

Total aerobic microbial count (cfu/gm)

By plate method when incubated at 30-35°C for not less than 3 days.

Bacterial Count : <= 2000 CFU/gram

Total Yeast and mould count (cfu/gm)

By plate method when incubated at 20-25°C for not less than 5 days.

Yeast & mould Count : <= 100 CFU/gram

Test for Pathogens

1. *Escherichia coli*-Negative in 10 gms of sample 2. *Salmonella* species-Negative in 10 gms of sample 3. *Pseudomonas aeruginosa*- Negative in 10 gms of sample 4. *Staphylococcus aureus*- Negative in 10 gms of sample 5. *Candida albicans*- Negative in 10 gms of sample 6. *Clostridia*- Negative in 10 gms of sample

Indole test

Tryptophan content: Passes

Cultural response

Cultural response observed after incubation at 35 - 37°C for 18-48 hours by preparing HiVeg™ Soyabean Casein Digest Medium (MV011), using HiVeg™ Hydrolysate as an ingredient.

Cultural Response

Organism	Growth
<i>Escherichia coli</i> ATCC 25922	Characteristic, luxuriant growth
<i>Pseudomonas aeruginosa</i> ATCC 27853	Characteristic, luxuriant growth
<i>Enterobacter aerogenes</i> ATCC 13048	Characteristic, luxuriant growth
<i>Salmonella</i> Typhi ATCC 6539	Characteristic, luxuriant growth
<i>Staphylococcus aureus</i> ATCC 25923	Characteristic, luxuriant growth

<i>Streptomyces albus</i> ATCC 3004	Characteristic, luxuriant growth
<i>Streptococcus pyogenes</i> ATCC 19615	Luxuriant w/beta haemolysis (with addition of sterile 5% sheep blood in above medium after 48 hours of incubation at 35-37°C).
<i>Neisseria gonorrhoeae</i> ATCC 19424	Luxuriant w/beta haemolysis (with addition of sterile 10% sheep blood to above medium heated to 80 to 90°C until blood has turned to chocolate brown and incubated in 10% CO ₂ atmosphere after 48 hours of incubation at 35-37°C).

Chemical Analysis

Total Nitrogen	>= 11.0%
Amino Nitrogen	>= 3.0%
Sodium chloride	<= 5.0%
Loss on drying	<= 7.0%
Residue on ignition	<= 15.0%

Storage and Shelf Life

Store between 10- 30°C in tightly closed container and away from bright light. Use before expiry date on label. On opening, product should be properly stored in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use.



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.