



## HiVeg™ Special Peptone

RM015V

It can be used for the preparation of media for cultivation of following bacteria: Neisseria species : G C HiVeg Agar Base (MV434), Thayer Martin HiVeg Medium Base (MV413) Yersinia species : Yersinia Selective HiVeg Agar Base (MV843). Staphylococci and Streptococci : Columbia Blood Agar Base HiVeg (MV144) . It can also be used for cultivation of other fastidious bacteria on large scale.

### Principle And Interpretation

HiVeg Special Peptone is manufactured under controlled conditions from vegetable proteins. It is especially adapted for the preparation of media for culturing fastidious bacteria and supports cultural characteristics comparable with Peptone special (RM015).

### Quality Control

#### Appearance

Light yellow to yellow, may have a slight green tinge ,homogenous free flowing powder, having characteristic odour of protein, derived from vegetable source.

#### Solubility

Freely soluble in distilled/ purified water, insoluble in alcohol.

#### Clarity

1% w/v aqueous solution is clear without any haziness after autoclaving at 15 lbs pressure (121°C ) for 15 minutes.

#### Reaction

Reaction of 2% w/v aqueous solution at 25°C.

#### pH

5.50- 7.50

#### Microbial Load:

##### Total aerobic microbial count (cfu/gm)

By plate method when incubated at 30-35°C for not less than 3 days.

Bacterial Count : <= 2000 CFU/gram

##### Total Yeast and mould count (cfu/gm)

By plate method when incubated at 20-25°C for not less than 5 days.

Yeast & mould Count : <= 100 CFU/gram

#### Test for Pathogens

1. *Escherichia coli*-Negative in 10 gms of sample 2. *Salmonella* species-Negative in 10 gms of sample 3. *Pseudomonas aeruginosa*- Negative in 10 gms of sample 4. *Staphylococcus aureus*- Negative in 10 gms of sample 5. *Candida albicans*- Negative in 10 gms of sample 6. *Clostridia*- Negative in 10 gms of sample

#### Indole test

Tryptophan content: Passes

#### Cultural response

Cultural response observed after incubation at 35 - 37°C for 18-48 hours by preparing Columbia Blood Agar Base HiVeg (MV144) using HiVeg™ Special Peptone as an ingredient.

#### Cultural Response

Organism	Growth	Haemolysis
<b>Cultural response</b>		
<i>Neisseria meningitidis</i> ATCC 13090	Luxuriant	none
<i>Staphylococcus aureus</i> ATCC 25923	Luxuriant	beta / gamma
<i>Staphylococcus epidermidis</i> ATCC 12228	Luxuriant	gamma
<i>Streptococcus pneumoniae</i> ATCC 6303	Luxuriant	alpha

---

*Streptococcus pyogenes* Luxuriant beta  
ATCC 19615

### Chemical Analysis

Total Nitrogen	>= 11.50%
Amino Nitrogen	>= 3.80%
Sodium chloride	<= 4.0%
Loss on drying	<= 7.0%
Residue on ignition	<= 15.0%

### Storage and Shelf Life

Store below 30°C. Use before expiry date on the label.



#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.