



Granulated Media

Reduce Dust Generation &
Allergic Responses



WHAT IF YOUR CHOICE OF MEDIA REDUCES DUST GENERATION AND ALLERGIC RESPONSES...

Media components in powdered media may lead to allergic responses if inhaled by personnel. Our unique granulation technology converts traditional powdered media into strong multi-particles. Granulated Media combines the advantage of safety with the ease of media preparation.



Granulated Media

- Control of media aerosols
- No lump formation
- Superior flow properties
- Higher bulk density
- Enables faster dissolution



HiMedia's granulated culture media, combines the high throughput technology of granulation and production of dehydrated culture media. The effective media has similar qualitative attributes to that of powdered dehydrated culture media with several added benefits in physical parameters.

Uniform blends of culture media

Individual components of dehydrated culture media tend to segregate in transport or handling because of the difference in their particle size, shape, and density. Granulated media from HiMedia forms uniform homogenous consistency, which eliminates segregation and facilitates consistent analysis.

Superior flow properties

The flow rate of granular media is better than powder media and it could also prevent bridging and caking issues associated with powder media. Most of the culture media components are hygroscopic and granulation tends to minimize this issue.

Control of media aerosols

Dust/Powder is generally difficult to handle in culture media preparation and may pose health-related issues to the microbiologist upon exposure to the media aerosols. Antibiotics and several other selective agents used in various media preparations may result in allergic responses and several other health issues in various parts of the body. These are effectively controlled in granulated form.

Higher bulk density

Increased bulk density has been shown to have improved the packaging, storage, and transport of dehydrated culture media.



Product	Code	Packing
A-1 Broth, Granulated for determining the presence of faecal coliforms in water samples and food by MPN technique. Gms/Lit : 31.50 15.87 Lit/500G	GM874-500G	500gm
APT Agar, Granulated for cultivation of heterofermentative <i>Lactobacilli</i> and other organisms requiring a high thiamine content. Gms/Lit : 61.18 8.17 Lit/500G	GM226-500G	500gm
Aero Pseudo Selective Agar, Granulated used for detecting <i>Pseudomonas</i> and <i>Aeromonas</i> in foodstuffs, waste water as well as equipment from the food industry. Gms/Lit : 44.86 11.15 Lit/500G Penicillin G sodium salt (100,000 IU), Pimaricin - 0.01 g/Lit ➤	GM1620-500G	500gm
Alicyclobacillus Agar, Granulated (BAT Agar, Granulated) used for isolation of <i>Alicyclobacillus</i> in fruit juice. Gms/Lit : 28.95 17.27 Lit/500G	GM1650-500G	500gm
Alkaline Peptone Water, Granulated for enrichment of <i>Vibrio</i> species from food, water and clinical samples. Gms/Lit : 20.00 25 Lit/500G	GM618-500G	500gm
Alternative Thioglycollate Medium, Granulated (Thioglycollate Broth, Alternative, Granulated) recommended for sterility testing of turbid or viscous biological products and for cultivation of anaerobes and microaerophiles from clinical and nonclinical specimens. Gms/Lit : 29.00 17.24 Lit/500G	GM010-500G	500gm
Anaerobic Agar (Brewer), Granulated for the isolation and sensitivity testing of anaerobic and microaerophilic organisms and study of colonial morphology. Gms/Lit : 53.00 9.43 Lit/500G	GM491-500G	500gm
Antibiotic Assay Medium No.11, Granulated (Erythromycin Seed Agar, Granulated) (Neomycin, Erythromycin Assay Agar, Granulated) for microbiological assay of antibiotics. (Note : pH of the Medium is 8.3 : ±0.2) Gms/Lit : 30.50 16.39 Lit/500G	GM004-500G	500gm
Antimicrobial Inhibitor Test Agar pH 6.0, Granulated for residual analysis of antimicrobial components in meat and organ samples using <i>Bacillus subtilis</i> (ATCC 6633) as test organism. Gms/Lit : 25.00 20 Lit/500G	GM1631-500G	500gm
Antimicrobial Inhibitor Test Agar pH 7.2, Granulated for residual analysis of antimicrobial components in meat and organ samples using <i>Bacillus subtilis</i> (ATCC 6633) as test organism. Gms/Lit : 25.34 19.73 Lit/500G	GM1601-500G	500gm







Product	Code	Packing
Antimicrobial Inhibitor Test Agar pH 8.0, Granulated for residual analysis of antimicrobial components in meat and organ samples using <i>Bacillus subtilis</i> (ATCC 6633) and <i>Micrococcus luteus</i> (ATCC 9341) as test organism. Gms/Lit : 26.07 19.18 Lit/500G	GM1632-500G	500gm
*Asparagine Broth, Granulated (Coccidioidin and Histoplasmin Broth, Granulated) for the preparation of Coccidioidin and Histoplasmin antigens for immunodiagnostic work. Gms/Lit : 28.01 17.85 Lit/500G	GM672-500G	500gm
Asparagine Proline Broth, Granulated for cultivation of <i>Pseudomonas aeruginosa</i> from water samples by membrane filter technique. Gms/Lit : 14.50 34.48 Lit/500G	GM1192-100G GM1192-500G	100gm 500gm
Azide Dextrose Broth, Granulated a selective medium for detection of <i>Streptococci</i> in water, sewage, food and other materials suspected of sewage contamination. Gms/Lit : 34.70 14.41 Lit/500G	GM345-500G	500gm
Azotobacter Broth (Glucose), Granulated for cultivation of Glucose positive <i>Azotobacter</i> species from soil. Gms/Lit : 16.4 30.49 Lit/500G	GM1721-500G	500gm
BAT Agar, Granulated (Alicyclobacillus Agar, Granulated) See : Alicyclobacillus Agar, Granulated	GM1650-500G	500gm
BHI Agar, Granulated (Special Infusion Agar, Granulated) for cultivation of fastidious pathogenic bacteria, yeasts and moulds. Gms/Lit : 52.00 9.62 Lit/500G Penicillin - 20 units/ml, Streptomycin - 40 µg/ml ➤	GM211-500G	500gm
BHI Broth, Granulated for propagation of pathogenic cocci and other fastidious organisms associated with blood culture work and allied pathological investigations. Gms/Lit : 37.00 13.51 Lit/500G	GM210-500G	500gm
BLE Broth Base, Granulated (Buffered Listeria Enrichment Broth Base, Granulated) recommended by FDA Committee for enrichment procedure for isolation of <i>Listeria monocytogenes</i> . Gms/Lit : 47.95 10.43 Lit/500G	GM1578-500G	500gm
*ANC Selective Supplement No. of Vials : 21 vials △	FD0631-5VL FD0631-5X5VL	5vl 5x5vl









Product	Code	Packing
Baird Parker Agar Base, Granulated for isolation and enumeration of coagulase positive Staphylococci from food and other materials. Gms/Lit : 63.00 7.94 Lit/500G	GM043-500G	500gm
*Egg Yolk Tel Emulsion (100 ml per vial) No. of Vials : 8 vials △ 4 vials △	FD046L-50MLX5VL FD046-100MLX5VL	50mlx5vl 100mlx5vl
*Egg Yolk Emulsion (50ml / 100 ml per vial) No. of Vials : 8 vials △ 4 vials △	FD045L-50MLX5VL FD045-100MLX5VL	50mlx5vl 100mlx5vl
*Pte 3.5% Selective Supplement (1 ml per vial) No. of Vials : 3 vials △	FD047-5VL	5vl
*BP S Selective Supplement (if desired) No. of Vials : 8 vials △	FD069-5VL	5vl
*FPT Inhibitor No. of Vials : 80 vials △	FD195-5VL FD195-5X5VL	5vl 5x5vl
Baird Parker Agar Base, Granulated recommended for the enumeration of coagulase positive Staphylococci from food and animal feeding stuffs. The composition and performance criteria are in accordance with ISO 6888-1:1999. Gms/Lit : 58.00 8.62 Lit/500G	GM0431-500G	500gm
*Egg Yolk Emulsion (50ml / 100 ml per vial) No. of Vials : 9 vials △	FD045L-50MLX5VL	50mlx5vl
*Pte 1% Selective Supplement (1 ml per vial) No. of Vials : 9 vials △	FD052-5VL FD052-5X5VL	5vl 5x5vl
*BP S Selective Supplement ➤ No. of Vials : 9 vials △	FD069-5VL	5vl
Baird Staphylococcus Enrichment Broth Base, Granulated for selective enrichment of pathogenic Staphylococci. Gms/Lit : 43.00 11.63 Lit/500G	GM1091-500G	500gm
*Pte 1% Selective Supplement (1 ml per vial) No. of Vials : 12 vials △	FD052-5VL FD052-5X5VL	5vl 5x5vl
Bi.G.G.Y. Agar, Granulated (Nickerson Medium, Granulated) for detection, selective isolation, differentiation and presumptive identification of <i>Candida albicans</i> and <i>Candida tropicalis</i> . Gms/Lit : 45.00 11.11 Lit/500G	GM217-500G	500gm
Bifidobacterium Selective Count Agar Base, Granulated (BSC Propionate Agar Base, Granulated) for enumeration of presumptive Bifidobacteria by colony count technique from milk products. Gms/Lit : 62.35 8.01 Lit/500G	GM1734-500G	500gm
*Mup Selective Supplement No. of Vials : 17 vials △	FD250-5VL	5vl
*1% glaci Selective Supplement No. of Vials : 17 vials △	FD251-5VL	5vl




Product	Code	Packing
Bile Esculin Azide Agar, Granulated for isolation and presumptive identification of faecal Streptococci. The composition and performance criteria of this medium are as per the specifications laid down in ISO 7899-2:2000. Gms/Lit : 56.65 8.83 Lit/500G	GM4931-500G	500gm
Bismuth Sulphite Agar, Granulated for selective isolation of Salmonellae from faeces, urine, sewage and other materials. Gms/Lit : 52.33 9.55 Lit/500G	GM027-500G	500gm
Blood Agar Base, Granulated (Infusion Agar, Granulated) recommended for isolation and cultivation of many fastidious pathogenic microorganisms like <i>Neisseria</i> , <i>Streptococci</i> after addition of blood. Gms/Lit : 40.00 12.5 Lit/500G	GM073-500G	500gm
Blood Agar Base No. 2 w/ 1.2% Agar, Granulated especially devised to permit the maximum recovery of fastidious pathogenic microorganisms without interfering with their haemolytic reactions. Gms/Lit : 39.50 12.66 Lit/500G	GM834A-500G	500gm
*NPBCVN Selective Supplement No. of Vials : 26 vials △	FD005-5VL FD005-5X5VL	5vl 5x5vl
*Blaser-Wang Selective Supplement No. of Vials : 26 vials △	FD006-5VL	5vl
*Butzler Selective Supplement No. of Vials : 26 vials △	FD007-5VL FD007-5X5VL	5vl 5x5vl
*Skirrow Selective Supplement No. of Vials : 26 vials △	FD008-5VL	5vl
*Minerals Growth Supplement No. of Vials : 26 vials △	FD009-5VL	5vl
*NNP Selective Supplement No. of Vials : 26 vials △	FD031-5VL FD031-5X5VL	5vl 5x5vl
Blood Free Campylobacter Selectivity Agar Base, Granulated for selective isolation and differentiation of <i>Campylobacter</i> species from food and animal feeding stuffs. The composition and performance criteria of this medium are as per the specifications laid down in ISO 10272-1&2:2017. Gms/Lit : 45.50 10.99 Lit/500G	GM887-500G	500gm
*CF Selective Supplement I No. of Vials : 22 vials △	FD067-5VL	5vl
*CAT Selective Supplement No. of Vials : 22 vials △	FD145-5VL	5vl
Bolton Broth Base, Granulated selective enrichment of <i>Campylobacter</i> species from food and animal feeding stuffs. The composition and performance criteria of this medium are as per the specifications laid down in ISO 10272-1:2017. Gms/Lit : 27.61 18.11 Lit/500G	GM1592-500G	500gm
*Bolton Selective Supplement No. of Vials : 37 vials △	FD231-5VL	5vl





Product	Code	Packing
Brilliant Green Agar Base w/ 1.2% Agar, Granulated recommended as enrichment medium for isolation of <i>Salmonellae</i> from faeces, urine and other pathological materials. Gms/Lit : 50.09 9.98 Lit/500G	GM016A-500G	500gm
*S Selective Supplement No. of Vials : 20 vials △	FD068-5VL	5vl
Brilliant Green Agar Base w/ Phosphates, Granulated for selective isolation of <i>Salmonellae</i> while inhibiting <i>Escherichia coli</i> , <i>Proteus</i> and <i>Pseudomonas</i> species. Gms/Lit : 51.69 9.67 Lit/500G	GM971-500G	500gm
*S Selective Supplement No. of Vials : 20 vials △	FD068-5VL	5vl
Bromo Cresol Purple Broth w/ Lactose, Granulated for identification of <i>Escherichia coli</i> and coliform bacteria from water samples. Gms/Lit : 28.02 17.84 Lit/500G	GM1265-500G	500gm
Brucella Agar Base, Granulated for selective isolation and cultivation of <i>Brucella</i> or <i>Campylobacter</i> species from clinical and non-clinical specimens. Gms/Lit : 43.10 11.6 Lit/500G	GM074-500G	500gm
*NPBCVN Selective Supplement No. of Vials : 24 vials △	FD005-5VL	5vl
	FD005-5X5VL	5x5vl
*Blaser-Wang Selective Supplement No. of Vials : 24 vials △	FD006-5VL	5vl
*Butzler Selective Supplement No. of Vials : 24 vials △	FD007-5VL	5vl
	FD007-5X5VL	5x5vl
*Skirrow Selective Supplement No. of Vials : 24 vials △	FD008-5VL	5vl
*Minerals Growth Supplement No. of Vials : 24 vials △	FD009-5VL	5vl
**Horse Serum No. of Vials : 600 ml △	RM1239-100ML	100ml
Bryant and Burkey Medium, Granulated for detecting and enumerating spores of lactate fermenting <i>Clostridium</i> in milk and dairy products. Gms/Lit : 33.00 15.15 Lit/500G	GM1385-500G	500gm
Buffered Glucose Broth, Granulated (Glucose Phosphate Broth, Granulated) (MR-VP Medium, Granulated) for studying Methyl Red and Voges Proskauer tests to differentiate amongst coli-aerogenes group Gms/Lit : 17.00 29.41 Lit/500G	GM070-500G	500gm
Buffered Glucose HiVeg™ Broth, Granulated (Glucose Phosphate HiVeg™ Broth, Granulated) (MR-VP HiVeg™ Medium, Granulated) for studying Methyl Red and Voges Proskauer tests to differentiate amongst coli-aerogenes group Gms/Lit : 17.00 29.41 Lit/500G	GMV070-500G ☉	500gm


Product	Code	Packing
Buffered Glucose Broth, Granulated (MR-VP Medium, Granulated) for studying Methyl Red and Voges Proskauer tests to differentiate amongst coli-aerogenes group. It is recommended by BIS committee under the specifications IS:5887 (Part I and Part V)-1976, Reaffirmed 2005. Gms/Lit : 15.00 33.33 Lit/500G	GM070S-500G	500gm
Buffered Listeria Enrichment Broth Base, Granulated (BLE Broth Base, Granulated) See : BLE Broth Base, Granulated	GM1578-500G	500gm
Buffered Peptone Water, Granulated for pre-enrichment medium used for increasing the recovery of injured <i>Salmonella</i> species from food prior to selective enrichment and isolation. Gms/Lit : 20.00 25 Lit/500G	GM614-500G	500gm
*CCV Supplement ➤ No. of Vials : 25 vials △	FD247-5VL	5vl
	FD247-5X5VL	5x5vl
Buffered Peptone Water, Granulated recommended as a pre-enrichment medium of <i>Enterobacteriaceae</i> organisms such as <i>Salmonella</i> and <i>Cronobacterium</i> species from food and animal feeding stuffs, water, milk, milk products and other products. The composition and performance criteria of this medium are as per the specifications laid down in ISO 6579-1:2017, ISO 21528-1:2004, ISO 22964:2017. Gms/Lit : 20.07 24.91 Lit/500G	GM1494I-500G	500gm
Buffered Peptone Water w/ NaCl, Granulated recommended as a diluent for carrying out microbial limit test from various specimens. Gms/Lit : 16.09 31.08 Lit/500G Polysorbate 20 or Polysorbate 80 - 0.1 to 1% ➤	GM1275-500G	500gm
Buffered Sodium Chloride-Peptone Solution, pH 7.0, Granulated ☹️ recommended as a diluent for carrying out microbial limit testing by harmonized methodology of pharmaceutical products in accordance with USP/EP/BP/JP/IP. Gms/Lit : 14.64 34.15 Lit/500G	GMH1275-500G	500gm
C.L.E.D. Agar w/ Bromo Thymol Blue, Granulated for isolation and differentiation of urinary pathogens on the basis of lactose fermentation. Gms/Lit : 36.15 13.83 Lit/500G	GM792-500G	500gm
Calcium M- Protein Agar, Granulated for detection and enumeration of proteolytic microorganisms in foodstuffs and other materials. Gms/Lit : 30.19 16.56 Lit/500G	GM1309-500G	500gm
Casein Soyabean Digest Agar, Granulated (Soyabean Casein Digest Agar, Granulated) (Tryptone Soya Agar, Granulated) a general purpose medium used for cultivation of a wide variety of microorganisms from clinical and non-clinical samples and for sterility testing in pharmaceutical products. Gms/Lit : 40.00 12.5 Lit/500G	GM290-500G	500gm
Casein-Soyabean Digest Agar, Granulated (Soybean-Casein Digest Agar, Granulated) ☹️ a general purpose medium used for cultivation of a wide variety of microorganisms from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP. Gms/Lit : 40.00 12.5 Lit/500G	GMH290-500G	500gm

Product	Code	Packing
Casein Soyabean Digest Broth, Granulated (Soyabean Casein Digest Medium, Granulated)  a general purpose medium used for cultivation of a wide variety of microorganisms and for sterility testing of moulds and lower bacteria in accordance with the harmonized method of USP/EP/BP/JP/IP Gms/Lit : 29.77 16.8 Lit/500G	GMH011-500G	500gm
Casein Soyabean Digest Broth, Granulated, Sterile (Soyabean Casein Digest Medium, Granulated, Sterile)  is γ irradiated sterile medium recommended for evaluation of sterility in manufacturing process. Gms/Lit : 29.77 16.8 Lit/500G	GMH011G-500G	500gm
Cetrimide Agar Base, Granulated for selective isolation of <i>Pseudomonas aeruginosa</i> from clinical specimens. Gms/Lit : 46.70+10ml glycerol 10.71 Lit/500G	GM024-500G	500gm
*NA Selective Supplement  No. of Vials : 11 vials 	FD130-5VL	5vl
Cetrimide Agar, Granulated  recommended as a selective medium used for the isolation of <i>Pseudomonas aeruginosa</i> from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP. Gms/Lit : 45.30+10ml glycerol 11.04 Lit/500G	GMH024-500G	500gm
China Blue Lactose Agar, Granulated for differentiation and enumeration of bacteria in milk. Gms/Lit : 38.30 13.05 Lit/500G	GM580-500G	500gm
*Coccidioidin and Histoplasmin Broth, Granulated (Asparagine Broth, Granulated) See : Asparagine Broth, Granulated	GM672-500G	500gm
Columbia Blood Agar Base, Granulated an efficient base for preparation of blood agar, chocolate agar and for various selective and identification media. Gms/Lit : 44.00 11.36 Lit/500G	GM144-500G	500gm
*NPBCVN Selective Supplement No. of Vials : 23 vials 	FD005-5VL FD005-5X5VL	5vl 5vl
*Blaser-Wang Selective Supplement No. of Vials : 23 vials 	FD006-5VL	5vl
*Butzler Selective Supplement No. of Vials : 23 vials 	FD007-5VL FD007-5X5VL	5vl 5x5vl
*Skirrow Selective Supplement No. of Vials : 23 vials 	FD008-5VL	5vl
*Minerals Growth Supplement No. of Vials : 23 vials 	FD009-5VL	5vl
*NC Selective Supplement No. of Vials : 23 vials 	FD030-5VL	5vl
*NNP Selective Supplement No. of Vials : 23 vials 	FD031-5VL FD031-5X5VL	5vl 5x5vl


Product	Code	Packing
*GNA Selective Supplement No. of Vials : 23 vials 	FD056-5VL	5vl
*VTCA Selective Supplement No. of Vials : 23 vials 	FD090-5VL	5vl
*Butzler VI Selective Supplement No. of Vials : 23 vials 	FD106-5VL FD106-5X5VL	5vl 5x5vl
*CO Selective Supplement No. of Vials : 23 vials 	FD119-5VL FD119-5X5VL	5vl 5x5vl
Columbia Agar, Granulated  for detection of <i>Clostridium sporogenes</i> from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP (Medium 15). Gms/Lit : 44.00 11.36 Lit/500G	GMH144-500G	500gm
*Gen Selective Supplement No. of Vials : 12 vials 	FD252-5VL	5vl
Czapek Dox Agar, Granulated a semisynthetic medium for general cultivation of fungi.	GM075-500G	500gm
Czapek Dox Agar, Modified, Granulated (Modified Czapek Dox Agar, Granulated) for cultivation and maintenance of fungi Gms/Lit : 45.36 11.02 Lit/500G	GM1170-500G	500gm
*10% Lactic acid solution (10 ml per vial) No. of Vials : 22 vials 	FD095-5VL FD095-5X5VL	5vl 5x5vl
Czapek Dox Broth, Granulated a semisynthetic medium used for cultivation of fungi. Gms/Lit : 35.01 14.28 Lit/500G	GM076-500G	500gm
DEV Lactose Peptone Broth, Granulated for enrichment and titre determination of coliform bacteria. Gms/Lit : 25.01 19.99 Lit/500G	GM1356-500G	500gm
DEV Tryptophan Broth, Granulated medium for subcultivation of coliform, differentiation and for indole testing in the bacteriological examination of water. Gms/Lit : 16.00 31.25 Lit/500G	GM1901-500G	500gm
D.T.M. Agar Base, Granulated (Dermatophyte Test Agar Base, Granulated) for selective isolation of dermatophytes. Gms/Lit : 40.20 12.44 Lit/500G	GM188-500G	500gm
*CCG Selective Supplement No. of Vials : 25 vials 	FD015-5VL	5vl
Deoxycholate Agar, Granulated for direct differential count of coliforms in dairy products and for isolation of enteric pathogens from rectal swabs, faeces and other pathological specimens. Gms/Lit : 45.03 11.1 Lit/500G	GM030-500G	500gm
Deoxycholate Citrate Agar, Granulated selective medium for the isolation of enteric pathogens particularly <i>Salmonella</i> and <i>Shigella</i> species. Gms/Lit : 70.52 7.09 Lit/500G	GM065-500G	500gm

Product	Code	Packing
Deoxycholate Citrate Agar Medium, Granulated for isolation of <i>Shigella</i> species from food samples. It is recommended by BIS committee under the specifications IS 5887 (Part 7):1999, Reaffirmed 2005. Gms/Lit : 55.45 9.02 Lit/500G	GM065S-500G	500gm
Deoxycholate Lactose Agar, Granulated for isolation and enumeration of coliforms in water, waste-water, milk and dairy products. Gms/Lit : 42.53 11.76 Lit/500G	GM066-500G	500gm
Dextrose Agar Base, Emmons, Granulated (Sabouraud Dextrose Agar Base, Modified, Granulated) for selective cultivation of pathogenic fungi. Gms/Lit : 47.00 10.64 Lit/500G	GM286-500G	500gm
*CC Supplement No. of Vials : 22 vials 	FD035-5VL	5vl
Dextrose Tryptone Agar, Granulated for detection and enumeration of mesophilic and thermophilic aerobic organisms in food. Gms/Lit : 30.04 16.64 Lit/500G	GM092-500G	500gm
▲ Dichloran Glycerol Medium Base, Granulated for selective isolation of xerophilic moulds from food samples. Gms/Lit : 31.60 15.82 Lit/500G	GM1129-500G	500gm
Dichloran Medium Base w/ Rose Bengal, Granulated for selective isolation and enumeration of yeasts and moulds associated with food spoilage Gms/Lit : 31.52 15.86 Lit/500G	GM1000-500G	500gm
*Chlor Selective Supplement No. of Vials : 32 vials 	FD033-5VL FD033-5X5VL	5vl 5x5vl
Differential Reinforced Clostridial Agar, Granulated for the enumeration and cultivation of Clostridia from water. Gms/Lit : 42.50 11.76 Lit/500G	GM1603-500G	500gm
Differential Reinforced Clostridial Broth Base, Granulated for the cultivation of Clostridia from water. Gms/Lit : 29.00 17.24 Lit/500G	GM549-500G	500gm
EC Broth, Granulated for selective enumeration of faecal & non-faecal coliform in water, waste water & shell fish Gms/Lit : 37.00 13.51 Lit/500G	GM127-500G	500gm
EE Broth, Mossel, Granulated for selective enrichment of <i>Enterobacteriaceae</i> in the bacteriological examination of food. Gms/Lit : 43.46 11.5 Lit/500G	GM287-500G	500gm
Enterobacteria Enrichment Broth Mossel, Granulated  for selective enrichment of <i>Enterobacteriaceae</i> from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP Gms/Lit : 42.93 11.65 Lit/500G	GMH287-500G	500gm

Product	Code	Packing
EMB Agar, Granulated for differential isolation of Gram-negative enteric bacilli from clinical and non-clinical specimens. Gms/Lit : 35.96 13.9 Lit/500G	GM317-500G	500gm
EMB Agar, Levine, Granulated for isolation, enumeration and differentiation of members of <i>Enterobacteriaceae</i> . Gms/Lit : 37.46 13.35 Lit/500G	GM022-500G	500gm
Endo Agar, Granulated for confirmation of the presumptive test for members of the coliform group. Gms/Lit : 41.50 12.05 Lit/500G	GM029-500G	500gm
Erythromycin Seed Agar, Granulated (Antibiotic Assay Medium No.11, Granulated) (Neomycin, Erythromycin Assay Agar, Granulated) See : Antibiotic Assay Medium No.11, Granulated	GM004-500G	500gm
Fluid Sabouraud Medium, Granulated (Sabouraud Medium, Fluid, Granulated) sterility test medium for moulds and lower bacteria in pharmaceutical preparations. Gms/Lit : 30.00 16.67 Lit/500G	GM013-500G	500gm
Fluid Selenite Cystine Medium, Granulated (Selenite Cystine Broth, Granulated) (Twin Pack) enrichment medium for isolation of <i>Salmonellae</i> in food, dairy products and materials of sanitary importance and clinical specimens. Gms/Lit : 19.01 gms of Part A + 4 gms of Part B  21.73 Lit/500G	GM025-500G	500gm
Fluid Tetrathionate Medium w/o Iodine and BG, Granulated (Tetrathionate Broth Base w/o Iodine and BG, Granulated) selective enrichment medium for isolating <i>Salmonellae</i> from food and other pathological materials. Gms/Lit : 46.00 10.87 Lit/500G Iodine solution - 20 ml/lit 0.1% Brilliant green solution - 10 ml/lit 	GM032-500G	500gm
Fluid Thioglycollate Medium, Granulated (Thioglycollate Medium, Fluid, Granulated) for sterility testing of biologicals and for cultivation of aerobes, anaerobes and microaerophiles. Gms/Lit : 29.75 16.81 Lit/500G	GM009-500G	500gm
Fraser Broth Base, Granulated recommended as a primary as well as secondary enrichment medium, for the isolation and enumeration of <i>Listeria monocytogenes</i> from food and animal feeds. Gms/Lit : 54.92 9.1 Lit/500G	GM1327-500G	500gm
*Fraser Selective Supplement No. of Vials : 9 vials 	FD1251-5VL	5vl
#Fraser Supplement No. of Vials : 18 vials 	FD141-5VL	5vl

 Sodium biselenite is also available in bud (DB001) and disc form (DD056). # On receipt store between 10-30°C.

* On receipt store between 2 - 8°C.  Harmonized Media  To be added but not provided.  On receipt store between 15-25°C

 Approx. number of vials required per 500gm of granulated medium.

Bulk supply of 2.5 kg, 5 kg etc is available on request.

Product	Code	Packing
*Fraser Broth w/Supplement, Granulated for the selective enrichment of <i>Listeria species</i> from food samples. Gms/Lit : 55.47 9.01 Lit/500G	GM2002-500G	500gm
Fraser Secondary Enrichment Broth Base, Granulated for the isolation, cultivation and enrichment of <i>Listeria monocytogenes</i> from food and environmental specimens. Gms/Lit : 57.85 8.64 Lit/500G	GM1083-500G	500gm
*Fraser Enrichment Supplement No. of Vials : 9 vials ▲	FD065-5VL	5vl
*Fraser Selective Supplement No. of Vials : 9 vials ▲	FD125-5VL	5vl
GN Broth, Hajna, Granulated for selective enrichment of Gram-negative enteric organisms. Gms/Lit : 39.00 12.82 Lit/500G	GM242-500G	500gm
Gassner Lactose Agar, Granulated for detection and isolation of pathogenic <i>Enterobacteriaceae</i> from food stuffs and other materials. Gms/Lit : 76.87 6.5 Lit/500G	GM1022-500G	500gm
Gelatin DEV Agar, Granulated for determining the total microbial count and detecting gelatin liquefying microorganisms in water. Gms/Lit : 50.00 10 Lit/500G	GM1609-500G	500gm
Giolitti-Cantoni Broth Base, Granulated for selective enrichment of <i>Staphylococcus aureus</i> from food. Gms/Lit : 54.20 9.23 Lit/500G	GM584-500G	500gm
*PTe 3.5% Selective Supplement (1 ml per vial) No. of Vials : 15 vials ▲	FD047-5VL	5vl
Glucose Phosphate Broth, Granulated (Buffered Glucose Broth, Granulated) (MR-VP Medium, Granulated) See : Buffered Glucose Broth, Granulated	GM070-500G	500gm
Glucose Phosphate HiVeg™ Broth, Granulated (Buffered Glucose HiVeg™ Broth, Granulated) (MR-VP HiVeg™ Medium, Granulated) See : Buffered Glucose HiVeg™ Broth, Granulated	GMV070-500G ●	500gm
Hanahan's Broth, Granulated (SOB Medium, Granulated) for use in cultivation of recombinant strains of <i>Escherichia coli</i> . Gms/Lit : 28.08 17.81 Lit/500G	GM1252-500G	500gm
Hektoen Enteric Agar, Granulated for differential and selective isolation of <i>Salmonella</i> and <i>Shigella</i> species from enteric pathological specimens. Gms/Lit : 76.67 6.52 Lit/500G	GM467-500G	500gm
Infusion Agar, Granulated (Blood Agar Base, Granulated) See : Blood Agar Base, Granulated	GM073-500G	500gm

Product	Code	Packing
Iron Sulphite Agar, Granulated for detection of thermophilic anaerobic organisms causing sulphide spoilage in food. Gms/Lit : 26.00 19.23 Lit/500G	GM868-500G	500gm
Jensen's Broth, Granulated for detection and cultivation of nitrogen fixing bacteria. Gms/Lit : 24.10 20.75 Lit/500G	GM973-500G	500gm
KF Streptococcus Agar Base w/ BCP, Granulated for detection and enumeration of faecal Streptococci. Gms/Lit : 76.41 6.54 Lit/500G	GM1007-500G	500gm
*TTC Solution 1% (10 ml per vial) No. of Vials : 7 vials ▲	FD057-5VL FD057-5X5VL	5vl 5x5vl
*Kimmig Fungi Agar Base, Granulated for cultivation, isolation and identification of fungi. Gms/Lit : 50.40 9.92 Lit/500G	GM1232-500G	500gm
*Kimmig Selective Supplement (Twin Pack) No. of Vials : 20 vials ▲	FD111-5VL	5vl
*George Kimmig Selective Supplement No. of Vials : 20 vials ▲	FD112-5VL	5vl
King's Medium A Base, Granulated for the non selective isolation, cultivation and pigment production of <i>Pseudomonas</i> species. Gms/Lit : 46.64 10.72 Lit/500G Glycerol - 10ml/lit ◀	GM1543-500G	500gm
King's Medium B Base, Granulated for the non selective isolation, cultivation and pigment production of <i>Pseudomonas</i> species. Gms/Lit : 42.23 11.84 Lit/500G Glycerol - 15ml/lit ◀	GM1544-500G	500gm
Kligler Iron Agar, Granulated for differential identification of Gram-negative enteric bacilli on the basis of the fermentation of dextrose, lactose and H ₂ S production. Gms/Lit : 57.52 8.69 Lit/500G	GM078-500G	500gm
Kundrat Agar, Modified, Granulated for the qualitative detection of residues of antibiotics and other chemotherapeutic agents in animal derived food. Gms/Lit : 40.52 12.34 Lit/500G	GM1360A-500G	500gm
LM Agar, Granulated (Liver Meat Agar, Granulated) for cultivation of fastidious anaerobes. Gms/Lit : 34.20 14.62 Lit/500G	GM1001-500G	500gm
Lactobacillus MRS Agar, Granulated (MRS Agar, Granulated) for isolation and cultivation of Lactobacilli. Gms/Lit : 67.15 7.45 Lit/500G	GM641-500G	500gm

◀ To be added but not provided. * On receipt store between 2 - 8°C.

▲ Approx. number of vials required per 500gm of granulated medium.

● The Animal based peptones in this medium have been fully replaced by TSE/BSE and GMO free Vegetable peptones.

Bulk supply of 2.5 kg, 5 kg etc is available on request.

Product	Code	Packing
*Lactobacillus MRS Agar (MRS agar), Granulated for the isolation and enumeration of lactic acid bacteria from meat and meat products. The composition and performance criteria of this medium are as per the specifications laid down in ISO 13721:1995. Gms/Lit : 65.13 7.68 Lit/500G	GM641I-500G	500gm
*Lactobacillus MRS Broth, Granulated (MRS Broth, Granulated) for cultivation of all Lactobacilli. Gms/Lit : 55.15 9.07 Lit/500G	GM369-500G	500gm
Lactose Broth, Granulated for the detection of <i>Enterobacteriaceae</i> particularly coliform and <i>Salmonella</i> species from water, food and dairy products as per Standard Methods of APHA and FDA BAM. Gms/Lit : 13.00 38.46 Lit/500G	GM1003-500G	500gm
Lauryl Sulphate Broth, Granulated (Lauryl Tryptose Broth, Granulated) for detection and enumeration of coliform bacteria in water, wastewater, dairy products and other food samples. Gms/Lit : 35.60 14.04 Lit/500G	GM080-500G	500gm
Legionella Agar Base, Granulated with addition of supplements it is used for cultivation of <i>Legionella</i> species. Gms/Lit : 25.00 20 Lit/500G	GM809A-500G	500gm
*BCYE Growth Supplement No. of Vials : 40 vials Δ	FD142-5VL	5vl
OR		
*BCYE Growth Supplement No. of Vials : 200 vials Δ	FD142X-5VL	5vl
	FD142X-5X5VL	5x5vl
*GVPC Selective Supplement No. of Vials : 40 vials Δ	FD143-5VL	5vl
	FD143-5X5VL	5x5vl
*BMPA Selective Supplement \blacktriangleleft No. of Vials : 40 vials Δ	FD144-5VL	5vl
	FD144-5X5VL	5x5vl
*Legi Growth Supplement w/o L-Cysteine No. of Vials : 40 vials Δ	FD206-5VL	5vl
	FD206-5X5VL	5x5vl
*GVPN Selective Supplement No. of Vials : 40 vials Δ	FD242-5VL	5vl
Leifson Agar, Granulated for isolation of <i>Salmonella</i> and <i>Shigella</i> species. Gms/Lit : 47.42 10.54 Lit/500G	GM1380-500G	500gm
Letheen Agar, Modified, Granulated (Modified Letheen Agar, Granulated) for screening cosmetic products for microbial contamination. Gms/Lit : 54.10 9.24 Lit/500G	GM946-500G	500gm
Letheen Broth, Modified, Granulated (Modified Letheen Broth, Granulated) for screening cosmetic products for microbial contamination. Gms/Lit : 42.80 11.68 Lit/500G	GM976-500G	500gm

Product	Code	Packing
Linden Grain Medium, Granulated for Media Fill process simulation for beverage bottling, to test for low acid beverage spoiling bacteria. Gms/Lit : 29.50 16.95 Lit/500G	GM1916-500G	500gm
Listeria Identification Agar Base, (PALCAM), Granulated for selective isolation and identification of <i>Listeria</i> species. Gms/Lit : 68.88 7.26 Lit/500G	GM1064-500G	500gm
*PALCAM Selective Supplement No. of Vials : 15 vials Δ	FD061-5VL	5vl
	FD061-5X5VL	5x5vl
Listeria Identification Broth Base, (PALCAM), Granulated for selective enrichment and identification of <i>Listeria</i> species. Gms/Lit : 47.38 10.55 Lit/500G	GM1090-500G	500gm
*PALCAM Selective Supplement No. of Vials : 22 vials Δ	FD061-5VL	5vl
	FD061-5X5VL	5x5vl
Listeria Oxford Medium Base, Granulated for isolation of <i>Listeria</i> species from pathological specimens. Gms/Lit : 55.50 9.01 Lit/500G	GM1145-500G	500gm
*Oxford Listeria Supplement No. of Vials : 18 vials Δ	FD071-5VL	5vl
*CM Selective Supplement No. of Vials : 18 vials Δ	FD126-5VL	5vl
Listeria Selective Broth Base, Granulated for selective isolation and cultivation of <i>Listeria monocytogenes</i> from clinical specimens. Gms/Lit : 36.00 13.89 Lit/500G	GM889-500G	500gm
*HiANC Selective Supplement No. of Vials : 14 vials Δ	FD063-5VL	5vl
*ANC Selective Supplement No. of Vials : 28 vials Δ	FD063I-5VL	5vl
	FD063I-5X5VL	5x5vl
*Listeria Selective Enrichment Broth, Granulated for selective enrichment of <i>Listeria</i> species in accordance with FDA/IDF-FIL Gms/Lit : 36.10 13.85 Lit/500G	GM1865-500G	500gm
Lowenstein Jensen Medium Base, Granulated (L.J. Medium, Granulated) for isolation and cultivation of <i>Mycobacterium</i> species from clinical samples. Gms/Lit : 37.24 8 Lit/500G Egg emulsion base - 1000 ml \blacktriangleleft Glycerol - 12 ml/lit \blacktriangleright	GM162-500G	500gm
*Gruft Selective Supplement \blacktriangleright No. of Vials : 34 vials Δ	FD053-5VL	5vl
Luria Agar, Granulated for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 35.00 14.29 Lit/500G	GM557-500G	500gm

Product	Code	Packing
Luria Agar Base, Miller's Modification, Granulated for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> with or without addition of glucose. Gms/Lit : 30.50 16.39 Lit/500G 20% glucose solution - 10 ml/lit ➤	GM1726-500G	500gm
Luria Bertani Agar, Miller, Granulated (Miller Luria Bertani Agar, Granulated) for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> for genetic and molecular biology studies; may be used for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 40.00 12.5 Lit/500G	GM1151-500G	500gm
Luria Bertani HiVeg™ Agar, Miller, Granulated (Miller Luria Bertani HiVeg™ Agar, Granulated) for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> for genetic and molecular biology studies; may be used for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 40.00 12.5 Lit/500G	GMV1151-500G ●	500gm
Luria Bertani Agar, Modified, Granulated for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> for genetic and molecular biology studies; may be used for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 30.00 16.67 Lit/500G	GM1151F-500G	500gm
Luria Bertani Agar, Miller pH 7.0, Granulated for the cultivation and maintenance of recombinant strains of <i>E. coli</i> for genetic and molecular biology studies, may be used for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 40 12.5 Lit/500G	GM2120-500G	500gm
Luria Bertani Broth, Miller pH 7.0, Granulated for the cultivation and maintenance of recombinant strains of <i>E. coli</i> for genetic and molecular biology studies, may be used for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 25 20 Lit/500G	GM2119-500G	500gm
Luria Bertani Broth, Miller, Granulated (Miller Luria Bertani Broth, Granulated) for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> for genetic and molecular biology studies; may be used for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 25.00 20 Lit/500G	GM1245-500G	500gm
Luria Bertani HiVeg™ Broth, Miller, Granulated (Miller Luria Bertani HiVeg™ Broth, Granulated) for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> for genetic and molecular biology studies; may be used for routine cultivation and estimation of not particularly fastidious microorganisms. Gms/Lit : 25.00 20 Lit/500G	GMV1245-500G ●	500gm

Product	Code	Packing
Luria Broth, Granulated for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> . Gms/Lit : 20.00 25 Lit/500G	GM575-500G	500gm
Luria Broth Base, Miller's Modification, Granulated for the cultivation and maintenance of recombinant strains of <i>Escherichia coli</i> with or without addition of glucose. Gms/Lit : 15.50 32.26 Lit/500G	GM1725-500G	500gm
Lysine Iron Agar, Granulated for differentiation of enteric organisms especially <i>Salmonella</i> arizonae, based on their ability to decarboxylate or deaminate lysine and to form hydrogen sulphide (H ₂ S). The composition of this medium is in accordance with FDA BAM and APHA. Gms/Lit : 34.56 14.47 Lit/500G	GM377-500G	500gm
M-Endo Agar LES, Granulated for enumeration of coliforms in water using a two step membrane filtration technique. Gms/Lit : 51.05 9.79 Lit/500G 95% ethanol - 20 ml/lit ◀	GM1106-500G	500gm
M-FC Agar Base, Granulated for detection and enumeration of faecal coliforms using membrane filtration technique at higher temperature (44.5°C). Gms/Lit : 52.10 9.6 Lit/500G	GM1122-500G	500gm
#Rosolic Acid (0.1 gm/vl) No. of Vials : 10 vials △	FD058-5VL	5vl
M17 Agar Base, Granulated for cultivation of lactic Streptococci and plaque assay of lactic bacteriophages. Gms/Lit : 33.25 15.04 Lit/500G Disodium β-Glycerophosphate - 19 gm/Litre ◀	GM929-500G	500gm
M17 Broth, Granulated for cultivation of lactic Streptococci and their bacteriophages. Gms/Lit : 42.25 11.83 Lit/500G	GM1029-500G	500gm
MR-VP Medium, Granulated (Glucose Phosphate Broth, Granulated) (Buffered Glucose Broth, Granulated) See : Buffered Glucose Broth, Granulated	GM070-500G	500gm
MR-VP HiVeg™ Medium, Granulated (Glucose Phosphate HiVeg™ Broth, Granulated) (Buffered Glucose HiVeg™ Broth, Granulated) See : Buffered Glucose HiVeg™ Broth, Granulated	GMV070-500G ●	500gm
MR-VP Medium, Granulated (Buffered Glucose Broth, Granulated) See : Buffered Glucose Broth, Granulated	GM070S-500G	500gm
MR-VP Medium, Granulated recommended for the performance of Methyl Red and Voges-Proskauer tests used in the differentiation of coli-aerogenes group, <i>Cronobacter</i> spp. This composition is as per the specifications laid down in ISO 6579-1:2017, ISO 22964:2017(E), FDA BAM. Gms/Lit : 17.00 29.41 Lit/500G	GM070I-500G	500gm

➤ If required use ◀ To be added but not provided. # On receipt store between 10-30°C.

△ Approx. number of vials required per 500gm of granulated medium.

● The Animal based peptones in this medium have been fully replaced by TSE/BSE and GMO free Vegetable peptones.

Bulk supply of 2.5 kg, 5 kg etc is available on request.

Product	Code	Packing
*MRS Agar, Granulated (Lactobacillus Agar, Granulated) See : Lactobacillus Agar, Granulated	GM641-500G	500gm
* MRS Agar, Granulated (Lactobacillus MRS Agar, Granulated) See : Lactobacillus MRS Agar, Granulated	GM641I-500G	500gm
*MRS Broth, Granulated (Lactobacillus Broth, Granulated) See : Lactobacillus Broth, Granulated	GM369-500G	500gm
MYP Agar Base, Granulated (Phenol Red Egg Yolk Polymyxin Agar Base, Granulated) for isolation and identification of pathogenic Staphylococci and <i>Bacillus</i> species. Gms/Lit : 46.03 10.86 Lit/500G	GM636-500G	500gm
*PolyB Selective Supplement No. of Vials : 22 vials △	FD003-5VL FD003-5X5VL	5vl 5x5vl
*Egg Yolk Emulsion (50ml / 100 ml per vial) No. of Vials : 22 vials △ 11 vials △	FD045L-50MLX5VL FD045-100MLX5VL	50mlx5vl 100mlx5vl
MacConkey Agar w/o CV, NaCl w/ 0.5% Sodium Taurocholate, Granulated for cultivation and differentiation of enteric bacteria, restricting swarming of <i>Proteus</i> species. Gms/Lit : 55.04 9.08 Lit/500G	GM082-500G	500gm
MacConkey Agar w/o CV, NaCl w/0.5% Bile Salts, Granulated for cultivation and differentiation of enteric bacteria, restricting swarming of <i>Proteus</i> species from specimens such as urine which may contain large number of <i>Proteus</i> species as well as potentially pathogenic Gram-positive organisms. Gms/Lit : 47.07 10.62 Lit/500G	GM082A-500G	500gm
MacConkey Agar w/0.15% Bile Salts, CV and NaCl, Granulated for selective isolation and differentiation of coliform organisms and other enteric pathogens from clinical and non-clinical samples. Gms/Lit : 51.53 9.7 Lit/500G	GM081-500G	500gm
MacConkey Agar, Granulated 🔄 MacConkey Agar is recommended for selective isolation and differentiation of coliforms from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP. Gms/Lit : 49.53 10.09 Lit/500G	GMH081-500G	500gm
MacConkey Broth w/ Neutral Red, Granulated for selective enrichment and enumeration of coliforms from water and food samples. Gms/Lit : 40.07 12.48 Lit/500G	GM007-500G	500gm
MacConkey Broth Purple w/BCP, Granulated for presumptive identification of coliforms from variety of specimens such as water, milk and food etc. Gms/Lit : 40.01 12.5 Lit/500G	GM083-500G	500gm
MacConkey Broth, Granulated 🔄 for the presumptive identification of coliforms from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP. Gms/Lit : 34.51 14.49 Lit/500G	GMH083-500G	500gm






Product	Code	Packing
Malachite Green Broth, Granulated for selective enrichment of <i>Pseudomonas aeruginosa</i> . Gms/Lit : 25.13 19.9 Lit/500G	GM1266-500G	500gm
Malt Extract Agar Base (w/ Mycological Peptone), Granulated for detection, isolation and enumeration of yeasts and moulds. Gms/Lit : 50.00 10 Lit/500G	GM137-500G	500gm
*10% Lactic acid solution (10 ml per vial) No. of Vials : 20 vials △	FD095-5VL FD095-5X5VL	5vl 5x5vl
Malt Extract Broth Base, Granulated for detection and enumeration of yeasts, moulds and aciduric microorganisms. Gms/Lit : 20.00 25 Lit/500G	GM255-500G	500gm
*10% Lactic acid solution (10 ml per vial) No. of Vials : 50 vials △	FD095-5VL FD095-5X5VL	5vl 5x5vl
Mannitol Salt Agar Base, Granulated for selective isolation of pathogenic Staphylococci. Gms/Lit : 111.02 4.50 Lit/500G	GM118-500G	500gm
*Egg Yolk Emulsion (50ml / 100 ml per vial) No. of Vials : 5 vials △ 3 vials △	FD045L-50MLX5VL FD045-100MLX5VL	50mlx5vl 100mlx5vl
Mannitol Salt Agar, Granulated 🔄 for selective isolation of pathogenic Staphylococci from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP. Gms/Lit : 111.02 4.50 Lit/500G	GMH118-500G	500gm
Maximum Recovery Diluent, Granulated a protective and isotonic diluent used for maximal recovery of microorganisms from a variety of sources. Gms/Lit : 9.50 52.63 Lit/500G	GM1030-500G	500gm
Microbial Content Test Agar, Granulated (Soyabean Casein Digest Agar w/Lecithin and Polysorbate 80, Granulated) (Tryptone Soya Agar w/ Lecithin & Polysorbate 80, Granulated) for determining efficiency of sanitization of containers, equipment surfaces, water miscible cosmetics etc. Gms/Lit : 45.70 10.94 Lit/500G	GM449-500G	500gm
Miller Luria Bertani Agar, Granulated (Luria Bertani Agar, Miller, Granulated) See : Luria Bertani Agar, Miller, Granulated	GM1151-500G	500gm
Miller Luria Bertani HiVeg™ Agar, Granulated (Luria Bertani HiVeg™ Agar, Miller, Granulated) See : Luria Bertani HiVeg™ Agar, Miller, Granulated	GMV1151-500G 🌱	500gm
Miller Luria Bertani Broth, Granulated (Luria Bertani Broth, Miller, Granulated) See : Luria Bertani Broth, Miller, Granulated	GM1245-500G	500gm




* On receipt store between 2 - 8°C.

△ Approx. number of vials required per 500gm of granulated medium. 🔄 Harmonized Media

🌱 The Animal based peptones in this medium have been fully replaced by TSE/BSE and GMO free Vegetable peptones.

Bulk supply of 2.5 kg, 5 kg etc is available on request.

Product	Code	Packing
Miller Luria Bertani HiVeg™ Broth, Granulated (Luria Bertani HiVeg™ Broth, Miller, Granulated) See : Luria Bertani HiVeg™ Broth, Miller, Granulated	GMV1245-500G 	500gm
Mineral Modified Glutamate Medium Base, Granulated (Double Strength) (Twin pack) for enumeration of coliform bacteria in water and waste water. Gms/Lit : 22.69 gms of Part A + 12.7 gms of Part B 14.13 Lit/500G	GM643-500G	500gm
Minerals Modified Glutamate Agar Base, Granulated (Twin Pack) for the enumeration of <i>Escherichia coli</i> from meat and meat products. The composition and performance criteria of this medium are as per the specifications laid down in ISO 6391:1997 & IS 15463:2004. Gms/Lit : 26.29 gms of Part A + 6.35 gms of Part B 15.32 Lit/500G Ammonium chloride - 2 gm/Litre 	GM643I-500G	500gm
Modified Czapek Dox Agar, Granulated (Czapek Dox Agar, Modified, Granulated) See : Czapek Dox Agar, Modified, Granulated	GM1170-500G	500gm
Modified EC Broth Base, Granulated for isolation of <i>Escherichia coli</i> O157 : H7 from food and other samples. Gms/Lit : 36.62 13.65 Lit/500G	GM1285-500G	500gm
*NO 5 Selective Supplement No. of Vials : 56 vials 	FD096-5VL FD096-5X5VL	5vl 5x5vl
Modified Lethen Agar, Granulated (Lethen Agar, Modified, Granulated) See : Lethen Agar, Modified, Granulated	GM946-500G	500gm
Modified Lethen Broth, Granulated (Lethen Broth, Modified, Granulated) See : Lethen Broth, Modified, Granulated	GM976-500G	500gm
Modified Rappaport Vassiliadis Medium, Granulated for selective enrichment of <i>Salmonellae</i> from food and environmental specimens. Gms/Lit : 30.07 16.63 Lit/500G	GM1137-500G	500gm
Modified Semisolid RV Medium Base, Granulated a semisolid medium for detection of motile <i>Salmonella</i> species from food and environmental specimens. Gms/Lit : 32.95 15.17 Lit/500G	GM1482-500G	500gm
*IMRV/RV Selective Supplement No. of Vials : 16 vials 	FD193-5VL	5vl
Modified Soyabean Bile Broth Base, Granulated recommended as an enrichment medium for the detection of <i>Escherichia coli</i> O157 : H7 from food and animal feeding stuffs. The composition and performance criteria are in accordance with ISO 16654:2001. Gms/Lit : 33.00 15.15 Lit/500G	GM1286I-500G	500gm
*NO 20 Selective Supplement No. of Vials : 16 vials 	FD290-5VL	5vl

Product	Code	Packing
Modified Tergitol 7 Agar Base, Granulated (Tergitol -7 Agar Base, Modified, Granulated) recommended for selective isolation and enumeration of coliform organisms in water by membrane filtration method. The composition and performance criteria are in accordance with ISO 9308-1:2014. Gms/Lit : 57.15 8.75 Lit/500G	GM616I-500G	500gm
*TTC Solution 1% (10 ml per vial) No. of Vials : 3 vials 	FD057-5VL FD057-5X5VL	5vl 5x5vl
Mueller Hinton Agar, Granulated for determination of susceptibility of microorganisms to antimicrobial agents. Gms/Lit : 38.00 13.16 Lit/500G	GM173-500G	500gm
Mueller Hinton Agar No. 2, Granulated for testing susceptibility of common and rapidly growing bacteria using antimicrobial discs by the Bauer-Kirby method, manufactured to contain low levels of thymine, thymidine, calcium and magnesium. Gms/Lit : 38.00 13.16 Lit/500G	GM1084-500G	500gm
Mueller Hinton Broth, Granulated recommended for determination of invitro susceptibility of bacterial strains against antibacterial agents by broth dilution methods. Gms/Lit : 21.00 23.81 Lit/500G	GM391-500G	500gm
Mueller Kauffman Tetrathionate Broth Base, Granulated for improved enrichment and isolation of <i>Salmonellae</i> . Gms/Lit : 82.05 6.09 Lit/500G	GM876-500G	500gm
Mueller Kauffman Tetrathionate Novobiocin Broth Base, Granulated for improved enrichment and isolation of <i>Salmonellae</i> as per ISO 6579-1:2017 Gms/Lit : 89.42 5.59 Lit/500G	GM1496I-500G	500gm
*MKTT Supplement No. of Vials : 6 vials  Iodine solution - 20 ml/litre 	FD203-5VL	5vl
Neomycin, Erythromycin Assay Agar, Granulated (Antibiotic Assay Medium No.11, Granulated) (Erythromycin Seed Agar, Granulated) See : Antibiotic Assay Medium No.11, Granulated	GM004-500G	500gm
Nickerson Medium, Granulated (Bi.G.G.Y. Agar, Granulated) See : Bi.G.G.Y. Agar, Granulated	GM217-500G	500gm
Nutrient Agar, Granulated for cultivation of less fastidious microorganisms, can be enriched with blood or other biological fluids. Gms/Lit : 28.00 17.86 Lit/500G	GM001-500G	500gm
Nutrient Agar No.2, Granulated used as a general purpose culture medium. Gms/Lit : 40.00 12.5 Lit/500G	GM1269-500G	500gm
Nutrient Agar, pH 7.0, Granulated for cultivation of <i>Salmonella</i> species. Gms/Lit : 23.00 21.74 Lit/500G	GM561A-500G	500gm

* On receipt store between 2 - 8°C.











 To be added but not provided.  Approx. number of vials required per 500gm of granulated medium.



 The Animal based peptones in this medium have been fully replaced by TSE/BSE and GMO free Vegetable peptones.






Bulk supply of 2.5 kg, 5 kg etc is available on request.









Product	Code	Packing
Nutrient Broth, Granulated for general cultivation of less fastidious microorganisms from clinical and non-clinical samples. It can be enriched with blood or other biological fluids. Gms/Lit : 13.00 38.46 Lit/500G	GM002-500G	500gm
Nutrient Broth, pH 6.9 w/o NaCl, Granulated recommended as general purpose medium for the cultivation of microorganisms from water samples. It is recommended by FDA BAM, APHA for enrichment of food samples detection of <i>Salmonella</i> . Gms/Lit : 8.00 62.5 Lit/500G	GM088-500G	500gm
OF Basal Medium, Granulated for differentiation of Gram-negative bacteria on the basis of fermentative and oxidative metabolism of carbohydrates. Gms/Lit : 9.38 53.3 Lit/500G 10% Dextrose, 10% Lactose, 10% Saccharose ◀	GM395-500G	500gm
Orange Serum Agar, Granulated for cultivation and enumeration of microorganisms associated with spoilage of citrus products, cultivation of Lactobacilli and other aciduric organisms and pathogenic fungi from food samples. Gms/Lit : 45.50 10.99 Lit/500G	GM933-500G	500gm
Oxytetra Glucose Yeast Agar Base, Granulated (OGYE Agar Base, Granulated) for selective isolation and enumeration of yeasts and/or moulds in food. Gms/Lit : 37.00 13.51 Lit/500G	GM639-500G	500gm
*OxyT Selective Supplement No. of Vials : 27 vials △	FD032-5VL FD032-5X5VL	5vl 5x5vl
Oxytetra Glucose Yeast Agar Base, Granulated (OGYE Agar Base, Granulated) for isolation and enumeration of yeasts & moulds from milk and milk products. The composition and performance criteria are in accordance with ISO 6611:2004. Gms/Lit : 37.00 13.51 Lit/500G	GM639I-500G	500gm
*OxyT Selective Supplement No. of Vials : 27 vials △	FD032-5VL FD032-5X5VL	5vl 5x5vl
*Genox Selective Supplement No. of Vials : 27 vials △	FD131-5VL FD131-5X5VL	5vl 5x5vl
PA Broth, Granulated for the detection of presence & absence of coliform bacteria in water from treatment plants or distribution systems. Gms/Lit : 30.51 16.39 Lit/500G	GM1186-500G	500gm
Peptone Granulated Granulated form of peptone recommended for general bacteriological work including mass cultivation.	RMG7709-500G	500gm
Peptone Type I, Bacteriological, Granulated used as culture media ingredient	RMG667-500G RMG667-5KG	500gm 5kg


Product	Code	Packing
Peptone water, Double buffered, Granulated recommended for the pre-enrichment of <i>Enterobacteriaceae</i> from foodstuffs and other test material. Gms/Lit : 25.00 20 Lit/500G	GM2124-500G	500gm
Perfringens Agar Base, Granulated (Tryptose Sulphite Cycloserine Agar Base, Granulated) (T.S.C./S.F.P. Agar Base, Granulated) with the addition of selective supplement and enrichment, it is used for the presumptive identification and enumeration of <i>Clostridium perfringens</i> . Gms/Lit : 47.00 10.64 Lit/500G	GM837-500G	500gm
*S.F.P. Selective Supplement No. of Vials : 22 vials △	FD013-5VL FD013-4X5VL	5vl 4x5vl
*T.S.C. Selective Supplement No. of Vials : 22 vials △	FD014-5VL FD014-4X5VL	5vl 4x5vl
*CMF Selective Supplement No. of Vials : 22 vials △	FD243-5VL	5vl
*Egg Yolk Emulsion (50ml / 100 ml per vial) No. of Vials : 11 vials △ 6 vials △	FD045L-50MLX5VL FD045-100MLX5VL	50mlx5vl 100mlx5vl
Phenol Red Agar Base, Granulated a basal medium to which carbohydrates may be added for use in fermentation studies of microorganisms. Gms/Lit : 31.02 16.12 Lit/500G	GM053-500G	500gm
Phenol Red Broth Base, Granulated a basal medium to which carbohydrates are added for determination of fermentation reactions of pure cultures of microorganisms. Gms/Lit : 16.02 31.21 Lit/500G	GM054-500G	500gm
Phenol Red Egg Yolk Polymyxin Agar Base, Granulated (MYP Agar Base, Granulated) See : MYP Agar Base, Granulated	GM636-500G	500gm
Pikovskaya's Broth (Medium), Granulated for detection of phosphate solubilizing microorganisms Gms/Lit : 16.30 30.67 Lit/500G	GM1719-500G	500gm
Plate Count Agar, Granulated (Standard Methods Agar, Granulated) for determination of plate counts of microorganisms in food, water, waste water. Gms/Lit : 23.50 21.28 Lit/500G	GM091-500G	500gm
Potato Dextrose Agar, Granulated for isolation and enumeration of yeasts and moulds from dairy and other food products. Gms/Lit : 39.00 12.82 Lit/500G 10% tartaric acid ➤	GM096-500G	500gm

Product	Code	Packing
Potato Dextrose Agar, Granulated  for the cultivation of yeasts and moulds from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP. Gms/Lit : 39.00 12.82 Lit/500G 10% tartaric acid 	GMH096-500G	500gm
Potato Dextrose Broth, Granulated for cultivation and enumeration of yeasts and moulds. Gms/Lit : 24.00 20.83 Lit/500G 10% tartaric acid 	GM403-100G GM403-500G	100gm 500gm
Pseudomonas Agar, Granulated (For Fluorescein) for detection of fluorescein production by <i>Pseudomonas</i> species. Gms/Lit : 38.00 13.16 Lit/500G Glycerol - 10 ml/lit 	GM120-500G	500gm
Pseudomonas Agar, Granulated (For Pyocyanin) for detection of pyocyanin production by <i>Pseudomonas</i> species. Gms/Lit : 46.40 10.78 Lit/500G Glycerol - 10 ml/lit 	GM119-500G	500gm
Pseudomonas Agar Base, Granulated for selective isolation of <i>Pseudomonas</i> species from environment, food, water samples. Gms/Lit : 48.40 10.33 Lit/500G Glycerol - 10 ml/lit 	GM085-500G	500gm
*Cetrinix Supplement No. of Vials : 21 vials 	FD029-5VL FD029-5X5VL	5vl 5x5vl
*CFC Supplement No. of Vials : 21 vials 	FD036-5VL FD036-5X5VL	5vl 5x5vl
R-2A Agar, Granulated for heterotrophic plate count of treated potable water using longer incubation periods. Gms/Lit : 18.12 27.59 Lit/500G	GM962-500G	500gm
Rappaport Vassiliadis Soya Broth, Granulated (RVS Broth, Granulated) recommended as a selective enrichment medium for the <i>Salmonellae</i> species, from the food and animal feeding stuffs and from clinical specimens. Gms/Lit : 27.11 18.44 Lit/500G	GM1491-500G	500gm
Rappaport Vassiliadis Salmonella Enrichment Broth, Granulated  for selective enrichment of <i>Salmonella</i> species from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP. Gms/Lit : 27.11 18.44 Lit/500G	GMH1491-500G	500gm
Reinforced Clostridial Agar, Granulated for the cultivation and enumeration of Clostridia and other anaerobes. Gms/Lit : 51.00 9.8 Lit/500G	GM154-500G	500gm
Reinforced Medium for Clostridia, Granulated  for the cultivation and enumeration of Clostridia from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP. Gms/Lit : 37.54 13.32 Lit/500G	GMH443-500G	500gm


Product	Code	Packing
Ringer Salt Solution, Granulated an isotonic diluent for food, milk and dairy products during microbiological examination. Gms/Lit : 8.91 11.22 Lit/100G	GM525-100G	100gm
Rinsing Fluid, Granulated used as a rinsing fluid in the membrane filtration procedure. Gms/Lit : 9.00 55.56 Lit/500G	GM1622-500G	500gm
*Rogosa SL Agar, Granulated recommended for selective cultivation of oral, vaginal and faecal Lactobacilli. Gms/Lit : 74.72 6.69 Lit/500G	GM130-500G	500gm
Rose Bengal Agar Base, Granulated for selective isolation and enumeration of yeasts and moulds from environmental materials, foodstuff and clinical samples. Gms/Lit : 31.55 15.85 Lit/500G	GM842-500G	500gm
*Chlor Selective Supplement No. of Vials : 32 vials 	FD033-5VL FD033-5X5VL	5vl 5x5vl
SIM Medium, Granulated for determination of hydrogen sulphide production, indole formation and motility of <i>enteric bacilli</i> . Gms/Lit : 36.23 13.8 Lit/500G	GM181-500G	500gm
SM Plate Count Agar, Granulated for determining the microbial count in milk and dairy products. Gms/Lit : 20.00 25 Lit/500G	GM1623-500G	500gm
SOB Medium, Granulated (Hanahan's Broth, Granulated) See : Hanahan's Broth, Granulated	GM1252-500G	500gm
SOC Broth, Granulated a medium for molecular biology. Gms/Lit : 28.08 17.81 Lit/500G	GM1379-500G	500gm
*SPS Agar, Granulated for detection of <i>Clostridium perfringens</i> in food samples. Gms/Lit : 40.03 12.49 Lit/500G	GM632-500G	500gm
SS Agar, Granulated (Salmonella Shigella Agar, Granulated) for differential and selective isolation of <i>Salmonella</i> and <i>Shigella</i> species from pathological specimens, suspected foodstuffs etc. Gms/Lit : 63.02 7.93 Lit/500G	GM108-500G	500gm
Sabouraud Dextrose Agar, Granulated for cultivation of yeasts, moulds and aciduric microorganisms. Gms/Lit : 65.00 7.69 Lit/500G	GM063-500G	500gm
Sabouraud Dextrose Agar, Granulated  for the cultivation of yeasts, moulds and aciduric bacteria from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP. Gms/Lit : 65.00 7.69 Lit/500G	GMH063-500G	500gm
Sabouraud Dextrose Agar Medium 4, Granulated for cultivation of yeasts, moulds and aciduric microorganisms in accordance with IP 2014. Gms/Lit : 61.36 8.15 Lit/500G	GMM063-500G	500gm


Product	Code	Packing
Sabouraud Dextrose Agar Base, Modified, Granulated (Dextrose Agar Base, Emmons, Granulated) See : Dextrose Agar Base, Emmons, Granulated	GM286-500G	500gm
Sabouraud Dextrose Broth, Granulated (Sabouraud Liquid Medium, Granulated) for cultivation of yeasts, moulds and aciduric microorganisms. Gms/Lit : 30.00 16.67 Lit/500G	GM033-500G	500gm
Sabouraud Dextrose HiVeg™ Broth, Granulated (Sabouraud Liquid HiVeg™ Medium, Granulated) for cultivation of yeasts, moulds and aciduric microorganisms. Gms/Lit : 30.00 16.67 Lit/500G	GMV033-500G 	500gm
Sabouraud Dextrose Broth, Granulated  for cultivation of yeasts, moulds and aciduric microorganisms from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/J.P. Gms/Lit : 30.00 16.67 Lit/500G	GMH033-500G	500gm
Sabouraud Dextrose Maltose Agar, Granulated for cultivation of moulds and yeasts and for testing antimycotic substances. Gms/Lit : 45.00 11.11 Lit/500G	GM1313-500G	500gm
Sabouraud Maltose Agar, Granulated for propagation of yeasts and moulds, particularly the parasitic fungi concerned with skin and scalp lesions. Gms/Lit : 65.00 7.69 Lit/500G	GM062-500G	500gm
Sabouraud Medium, Fluid, Granulated (Fluid Sabouraud Medium, Granulated) See : Fluid Sabouraud Medium, Granulated	GM013-500G	500gm
Sakazakii DHL Agar, Granulated used for the detection and isolation of pathogenic <i>Enterobacteriaceae</i> from all types of specimens. Gms/Lit : 63.71 7.85 Lit/500G	GM1619-500G	500gm
SalEnrich Broth Base, Granulated used for two-step enrichment of sublethally injured <i>Salmonellae</i> from food and feeds. Gms/Lit : 35.00 14.29 Lit/500G	GM1685-500G	500gm
#SalEnrich Selective Supplement No. of Vials : 58 vials 	FD238-5VL FD238-5X5VL	5vl 5x5vl
Salmonella Selective Enrichment Broth Base, Granulated for selective isolation and differentiation of <i>Salmonella</i> species. Gms/Lit : 25.00 20 Lit/500G	GM1843-500G	500gm
*NO 12 Selective Supplement No. of Vials : 20 vials 	FD275-5VL	5vl
Selenite Broth, Granulated (Selenite F Broth, Granulated) (Twin Pack) an enrichment medium for isolation of <i>Salmonella</i> species from faeces, urine or other pathological materials. Gms/Lit : 19.00 gms of Part A + 4 gms of Part B  21.74 Lit/500G	GM052-500G	500gm

Product	Code	Packing
Selenite Cystine Broth, Granulated (Fluid Selenite Cystine Medium, Granulated) (Twin Pack) See : Fluid Selenite Cystine Medium, Granulated	GM025-500G	500gm
Semisolid RV Medium Base, Granulated for isolation of <i>Salmonella</i> species from food stuffs and other materials based on selective motility. Gms/Lit : 30.20 16.56 Lit/500G	GM1428-500G	500gm
*IMRV/RV Selective Supplement No. of Vials : 34 vials 	FD193-5VL	5vl
Slanetz and Bartley Medium, Granulated for detection and enumeration of faecal Streptococci by membrane filtration technique. Gms/Lit : 46.50 10.75 Lit/500G	GM612-500G	500gm
Slanetz and Bartley Medium w/o TTC, Granulated for detection and enumeration of faecal Streptococci by membrane filtration technique. Gms/Lit : 46.40 10.78 Lit/500G	GM612A-500G	500gm
*TTC Solution 1% (10 ml per vial)  No. of Vials : 8 vials 	FD057-5VL FD057-5X5VL	5vl 5x5vl
Soyabean Casein Digest Agar, Granulated (Casein Soyabean Digest Agar, Granulated) (Tryptone Soya Agar, Granulated) See : Casein Soyabean Digest Agar, Granulated	GM290-500G	500gm
Soyabean Casein Digest Agar, Granulated  (Casein-Soyabean Digest Agar, Granulated) See : Casein-Soyabean Digest Agar, Granulated	GMH290-500G	500gm
Soyabean Casein Digest Agar w/ Lecithin and Polysorbate 80, Granulated (Microbial Content Test Agar, Granulated) (Tryptone Soya Agar w/ Lecithin & Polysorbate 80, Granulated) See : Microbial Content Test Agar, Granulated	GM449-500G	500gm
Soyabean Casein Digest Medium, Granulated (Tryptone Soya Broth, Granulated) a general purpose medium used for cultivation of a wide variety of microorganisms and sterility testing of moulds and lower bacteria as per various Pharmacopoeia. Gms/Lit : 30.00 16.67 Lit/500G	GM011-500G	500gm
Soyabean HiVeg™ Medium, Granulated (Tryptone Soya HiVeg™ Broth, Granulated)  a general purpose medium used for cultivation of a wide variety of microorganisms and sterility testing of moulds and lower bacteria as per various Pharmacopoeia. Gms/Lit : 30.00 16.67 Lit/500G	GMV011-500G 	500gm
Soyabean HiVeg™ Medium, Granulated, Sterile (Tryptone Soya HiVeg™ Broth, Granulated, Sterile)  is a γ-irradiated sterile powder recommended for the evaluation of sterility in manufacturing process. Gms/Lit : 30.00 16.67 Lit/500G	GMV011G-500G 	500gm




 Sodium biselenite is also available in bud (DB001) and disc form (DD056).





* On receipt store between 2 - 8°C.  Harmonized Media  If required use


 Approx. number of vials required per 500gm of granulated medium. # On receipt store between 10-30°C.



 The Animal based peptones in this medium have been fully replaced by TSE/BSE and GMO free Vegetable peptones.





Bulk supply of 2.5 kg, 5 kg etc is available on request.

Product	Code	Packing
Soyabean Casein Digest Medium, Granulated (Casein Soyabean Digest Broth, Granulated)  See : Casein Soyabean Digest Broth, Granulated	GMH011-500G	500gm
Soyabean Casein Digest Medium, Granulated, Sterile (Casein Soyabean Digest Broth, Granulated, Sterile)  See : Casein Soyabean Digest Broth, Granulated, Sterile	GMH011G-500G	500gm
Special Infusion Agar, Granulated (BHI Agar, Granulated) See : BHI Agar, Granulated	GM211-500G	500gm
Standard Methods Agar, Granulated (Plate Count Agar, Granulated) See : Plate Count Agar, Granulated	GM091-500G	500gm
Standard Nutrient Agar No. 1, Granulated for cultivation of fastidious bacteria. Gms/Lit : 37.00 13.51 Lit/500G	GM1210-500G	500gm
Standard Nutrient Agar No. 2, Granulated for the cultivation and enrichment of less fastidious bacteria. It can be used in detection of inhibitors in bacteriological examination of meat. Gms/Lit : 25.00 20 Lit/500G	GM1627-500G	500gm
Standard Nutrient Broth No. 1, Granulated for cultivation of fastidious bacteria. Gms/Lit : 25.00 20 Lit/500G	GM1224-500G	500gm
Sugar Free Agar, Granulated for examination of butter in accordance with International Dairy Federation. Gms/Lit : 34.00 14.71 Lit/500G	GM307-500G	500gm
Super Broth, Granulated for mass cultivation of <i>Escherichia coli</i> . Gms/Lit : 60.00 8.33 Lit/500G	GM1316-500G	500gm
Super Broth No.II, Granulated for cultivation of recombinant strains of <i>Escherichia coli</i> . Gms/Lit : 49.10 10.18 Lit/500G	GM1689-500G	500gm
T.A.T. Broth Base, Granulated for sterility testing of highly viscous or gelatinous substances such as salves, ointments and other cosmetic products. Gms/Lit : 25.00 20 Lit/500G Polysorbate 20 - 40ml/Lit 	GM562-500G	500gm
TCBS Agar, Granulated for selective isolation and cultivation of <i>Vibrio cholerae</i> and other enteropathogenic <i>Vibrio</i> 's causing food poisoning. Gms/Lit : 89.08 5.61 Lit/500G	GM189-500G	500gm
Tartoff - Hobbs Broth, Granulated (Terrific Broth, Granulated) for the cultivation of recombinant strains of <i>Escherichia coli</i> . Gms/Lit : 47.60 10.5 Lit/500G	GM1250-500G	500gm
Tergitol -7 Agar Base, Modified, Granulated (Modified Tergitol 7 Agar Base, Granulated) See : Modified Tergitol 7 Agar Base, Granulated	GM6161-500G	500gm

Product	Code	Packing
Terrific Broth, Granulated (Tartoff - Hobbs Broth, Granulated) See : Tartoff - Hobbs Broth, Granulated	GM1250-500G	500gm
Tetrathionate Brilliant Green Bile Broth, Granulated for isolation and identification of <i>Salmonellae</i> . Gms/Lit : 63.07 7.93 Lit/500G	GM1255-500G	500gm
Tetrathionate Broth Base w/o Iodine and BG, Granulated (Fluid Tetrathionate Medium w/o Iodine and BG, Granulated) See : Fluid Tetrathionate Medium w/o Iodine and BG, Granulated	GM032-500G	500gm
Thioglycollate Broth, Alternative, Granulated (Alternative Thioglycollate Medium, Granulated) See : Alternative Thioglycollate Medium, Granulated	GM010-500G	500gm
Thioglycollate Medium, Fluid, Granulated (Fluid Thioglycollate Medium, Granulated) See : Fluid Thioglycollate Medium, Granulated	GM009-500G	500gm
Triple Sugar Iron Agar, Granulated for identification of Gram-negative enteric bacilli on the basis of dextrose, lactose and sucrose fermentation and hydrogen sulphide production. Gms/Lit : 64.52 7.75 Lit/500G	GM021-500G	500gm
Tryptone Broth, Granulated (Tryptone Water, Granulated) for detection of indole producing microorganisms isolated from clinical and non-clinical samples. Gms/Lit : 15.00 33.33 Lit/500G	GM463-500G	500gm
Tryptone Glucose Extract Agar, Granulated (Tryptone Yeast Extract Agar, Granulated) for enumeration of bacteria in water, air, milk and dairy products. Gms/Lit : 24.00 20.83 Lit/500G	GM014-500G	500gm
Tryptone Soya Agar, Granulated (Casein Soyabean Digest Agar, Granulated) (Soyabean Casein Digest Agar, Granulated) See : Casein Soyabean Digest Agar, Granulated	GM290-500G	500gm
Tryptone Soya Agar w/ Lecithin & Polysorbate 80, Granulated (Microbial Content Test Agar, Granulated) (Soyabean Casein Digest Agar w/Lecithin and Polysorbate 80, Granulated) See : Microbial Content Test Agar, Granulated	GM449-500G	500gm
Tryptone Soya Broth, Granulated (Soyabean Casein Digest Medium, Granulated) See : Soyabean Casein Digest Medium, Granulated	GM011-500G	500gm
Tryptone Soya HiVeg™ Broth, Granulated (Soyabean HiVeg™ Medium, Granulated)  See : Soyabean HiVeg™ Medium, Granulated	GMV011-500G 	500gm
Tryptone Soya HiVeg™ Broth, Granulated, Sterile (Soyabean HiVeg™ Medium, Granulated, Sterile)  See : Soyabean HiVeg™ Medium, Granulated, Sterile	GMV011G-500G 	500gm

Product	Code	Packing
Tryptone Soya Yeast Extract Agar, Granulated for confirmation of <i>Listeria</i> in Henry's light. Gms/Lit : 51.00 9.8 Lit/500G	GM1214-500G	500gm
Tryptone Yeast extract Agar, Granulated for estimation of microbial counts in water. Gms/Lit : 21.00 23.81 Lit/500G	GM1272-500G	500gm
Tryptose Broth, Granulated for the cultivation primarily of <i>Brucella</i> species. Gms/Lit : 26.00 19.23 Lit/500G	GM177-500G	500gm
Tryptose Sulphite Cycloserine Agar Base, Granulated (Perfringens Agar Base, Granulated) (T.S.C./S.F.P. Agar Base, Granulated) See : Perfringens Agar Base, Granulated	GM837-500G	500gm
*Urea Agar Base, Granulated (Filter Sterilizable) (w/o Agar) with added agar it is used for detection of urea splitting microorganisms. Gms/Lit : 29.00 17.24 Lit/500G	GM112A-500G	500gm
*Urea Broth, Granulated (Filter Sterilizable) for identification of bacteria on the basis of urea utilization, specifically for the differentiation of <i>Proteus</i> species from <i>Salmonella</i> and <i>Shigella</i> species. Gms/Lit : 38.71 12.92 Lit/500G	GM111A-500G	500gm
Violet Red Bile Agar, Granulated for selective isolation, detection and enumeration of coli-aerogenes bacteria in water, milk other dairy food products and clinical samples. Gms/Lit : 41.53 12.04 Lit/500G	GM049-500G	500gm
Violet Red Bile Glucose Agar w/o Lactose, Granulated for enumeration of <i>Enterobacteriaceae</i> in raw food and clinical samples. The composition and performance criteria are in accordance with ISO 21528-1&2 :2017 Gms/Lit : 38.53 12.98 Lit/500G	GM581-500G	500gm
Violet Red Bile Glucose Agar, Granulated  for detection and enumeration of <i>Enterobacteriaceae</i> from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP. Gms/Lit : 40.62 12.31 Lit/500G	GMH581-500G	500gm
Wort Agar, Granulated for the cultivation and enumeration of yeasts. Gms/Lit : 48.28 10.36 Lit/500G	GM129-500G	500gm

Product	Code	Packing
Wort Broth, Granulated for cultivation and enumeration of yeasts. Gms/Lit : 33.28 15.02 Lit/500G Glycerol - 2.35 gm/Lit 	GM333-500G	500gm
2XYT Growth Medium, Granulated optimized formulation for the growth and maintenance of M13 phage or other filamentous ss DNA bacteriophages. Gms/Lit : 31.00 16.13 Lit/500G	GMG034-500G	500gm
Xylose Lysine Deoxycholate Agar, Granulated (XLD Agar, Granulated) for selective isolation and enumeration of <i>Salmonella</i> Typhi and other <i>Salmonella</i> species. Gms/Lit : 56.68 8.82 Lit/500G	GM031-500G	500gm
Xylose-Lysine Deoxycholate Agar, Granulated  a selective medium recommended for the isolation and enumeration of <i>Salmonella</i> species from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP Gms/Lit : 54.80 9.12 Lit/500G	GMH031-500G	500gm
YPD Broth, Granulated (YEPA Broth, Granulated) for the growth of <i>Saccharomyces cerevisiae</i> for molecular biology. Gms/Lit : 50.00 10 Lit/500G	GM1363-500G	500gm
YT Agar, Granulated for growth of <i>Escherichia coli</i> K12 strains used in preparation of phage and plasmid DNA according to Miller. Gms/Lit : 28.00 17.86 Lit/500G	GM1369-500G	500gm
YT Broth, Granulated (2X YT Broth, Granulated) for the cultivation of recombinant strains of <i>Escherichia coli</i> . Gms/Lit : 31.00 16.13 Lit/500G	GM1251-500G	500gm
Yeast extract Granulated Granulated form of peptone Yeast extract is manufactured from selected strains of <i>Saccharomyces cerevisiae</i> under control conditions by retaining all the nutritive values, amino acids, vitamins, especially B group and growth factors. It contains low salt and is recommended for microbiological media and for mass cultivation of various microorganisms.	RMG027-500G	500gm
▲ Yeast Glucose Chloramphenicol Agar, Granulated selective agar recommended for enumerating yeasts and moulds in milk and milk products. Gms/Lit : 40.10 12.47 Lit/500G	GM1590-500G	500gm



HiPetri Slim™ Plate

Salient features

- Widespread Industrial application (Water, Food & Cosmetics)
- Compact packing reduces storage space
- No Preparation time
- User-friendly, ready-to-use products
- Available in wide range of products that can be customised as per requirement

Eurasia

Africa

Australia

North America

Latin America

Europe

- CORPORATE OFFICE -

Plot No. C40, Road No. 21Y,
MIDC, Wagle Industrial Estate,
Thane (West) - 400604,
Maharashtra, India.

Tel : +91-22-6147 1919 /
6116 9797 / 6903 4800

Fax : +91-22-6147 1920

Email : info@himedialabs.com

OVERSEAS OFFICE

- North America -

HiMedia Laboratories LLC,
507 School House Rd,
Kennet Square 19348, PA, USA.

Tel : +1-484-734-4401

Fax : +1-484-734-4402

Email :

info@himedialabs.com

OVERSEAS OFFICE

- Europe -

HiMedia Laboratories GmbH,
Marie-Curie-Str. 3, 64683,
Einhausen, Germany.

Tel : +49 6251 989 24 26

Fax : +49 6251 989 24 27

Email :

infoeu@himedialabs.com

HiMedia is a global brand
with network reach
spanning over 150 countries
across the world.



OVERSEAS OFFICE

- Latin America -

HiMedia Laboratories
Rua Flórida, 1703 - Cj. 52
São Paulo, Brazil.

Tel : +55 11981141515

Email :

info@himedialabs.com

HiMedia Laboratories Pvt. Ltd.
www.himedialabs.com



HIMEDIA®

For Life is Precious



For Website