



Technical Datasheet

Penicillin Streptomycin Powder

With 5000 units Penicillin and 5mg Streptomycin per ml in 0.9% normal Saline when reconstituted with sterile tissue culture grade water to indicated volume

Product Code: A004

Product Description:

Penicillin and streptomycin are broad spectrum bacteriostatic and bacteriocidal agent effective against gram positive bacteria and gram negative bacteria. Penicillin acts by inhibiting cell wall synthesis. Streptomycin blocks the initiation complex of protein synthesis by binding to the 30S subunit of bacterial 70S ribosome, thus interfering with the protein synthesis.

A004 is a powdered antibiotic mixture formulated to contain 5000 units penicillin and 5mg streptomycin per ml when reconstituted with sterile tissue culture grade water to indicated level.

Directions:

- 1. Aseptically add 10ml sterile tissue culture grade water to one amber coloured bottle containing A004 powder.
- 2. Cap the bottle tightly and shake well to dissolve the powder.
- 3. Aseptically transfer this solution to one of the empty sterile bottle provided along with the product.
- 4. Again add 10ml sterile tissue culture grade water to the same amber coloured bottle of A004 to remove any traces of undissolved powder.
- 5. Add this solution to the bottle in which 10ml antibiotic solution was previously added (refer step 3).
- 6. Make up the volume to 50ml or 100ml (depending on pack size) by adding sterile tissue culture grade water up to the mark indicated on the bottle.

Recommended use concentration is 20ml/L. Sensitive cell lines may react differently to this product. Hence it is recommended to determine optimum usage dose empirically for individual cell line.

Quality Control:

Appearance

White homogeneous powder.

pН

4.00 - 7.00

Osmolality (mOsm/Kg H₂O)

305.00 -345.00

Penicillin Concentration

5000 U/ml

Streptomycin Concentration

5 mg/ml.

Antibiotic sensitivity test

Passes

Sterility

No bacterial or fungal growth is observed after 14 days of incubation, as per USP specification.

Cultural Response

No toxicity to cells.

Endotoxin Content

NMT 0.5 EU/ml

Storage and Shelf Life:

Store the powder at 2-8°C. Reconstituted solution should be stored at -30°C to -10°C.

Solution is stable for 3 days at 37°C and for approximately 2 to 3 weeks at 2 - 8°C.

Shelf life of the product is 24 months.

Use before expiry date given on the product label.

Disclaimer: Revision No.: 04/2025

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ Publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

