



Medium 199

With Hank's salts

Without L-Glutamine and Sodium bicarbonate

Product Code: AL192

Product Description:

Medium 199 was the first nutritionally defined medium developed by Morgan, Morton, and Parker in 1950. This complex medium was formulated specifically for nutritional studies on primary chick embryo fibroblasts in the absence of any additives. It was observed that explanted tissue could survive in Medium 199 without serum but long term cultivation of cells required supplementation of the medium with serum. Medium 199 is formulated with either Hank's salts or Earle's salts. The medium when supplemented with serum can be used for growth of a wide variety of cells. Medium 199 is presently used for the maintenance of non-transformed cells, vaccine and virus production and primary ex-plants of epithelial cells.

AL192 is 10X Medium 199 with Hank's salts. It does not contain L-glutamine and sodium bicarbonate. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

Composition:

Ingredients	mg/L
INORGANIC SALTS	
Calcium chloride dihydrate	1850.000
Ferric nitrate nonahydrate	7.200
Magnesium sulphate anhydrous	977.200
Potassium chloride	4000.000
Potassium phosphate monobasic	600.000
Sodium acetate anhydrous	500.000
Sodium chloride	80000.000
Sodium phosphate dibasic	478.600
AMINO ACIDS	
Glycine	500.000
L-Alanine	250.000
L-Arginine hydrochloride	700.000
L-Aspartic acid	300.000
L-Cysteine hydrochloride monohydrate	1.000
L-Cystine dihydrochloride	260.000
L-Glutamic Acid	670.000
L-Histidine hydrochloride monohydrate	220.000
L-Hydroxyproline	100.000

L-Isoleucine	200.000
L-Leucine	600.000
L-Lysine hydrochloride	700.000
L-Methionine	150.000
L-Phenylalanine	250.000
L-Proline	400.000
L-Serine	250.000
L-Threonine	300.000
L-Tryptophan	100.000
L-Tyrosine disodium salt	576.600
L-Valine	250.000
VITAMINS	
Ascorbic acid	0.5
Calciferol	1.000
Choline chloride	5.000
D-Biotin	0.1
D-Ca-Pantothenate	0.100
DL-Tocopherol phosphate disodium salt	0.100
Folic acid	0.100
Menadione	0.100
Nicotinamide	0.25
Nicotinic acid	0.25
Pyridoxal hydrochloride	0.25
Pyridoxine hydrochloride	1.400
Retinol Acetate	1.000
Riboflavin	1.000
Thiamine hydrochloride	0.100
i-Inositol	0.500
p-Amino benzoic acid (PABA)	0.500
OTHERS	
Adenine sulphate	100.000
Adenosine monophosphate	2.000
Adenosine triphosphate	10.000
Cholesterol	2.000
Deoxyribose	5.000
Glucose	10000.000
Glutathione reduced	0.500
Guanine hydrochloride	3.000
Hypoxanthine	3.54
Phenol red sodium salt	150.000
Polysorbate 80	49.000
Ribose	5.000

Thymine	3.000
Uracil	3.000
Xanthine	3.44

Directions:

1. Add 4.7ml of 7.5% sodium bicarbonate solution (TCL013) and 3.42ml of 200mM L-glutamine (TCL012) for 1 litre of 1X medium prior to use.

Material required but not provided:

L-Glutamine solution 200mM (TCL012)
Sodium bicarbonate solution 7.5% (TCL013)
Tissue culture grade water (TCL010)

Quality Control:**Appearance**

Red colored, clear solution

pH

4.20 -4.40

Sterility

No bacterial or fungal growth is observed after 14 days of incubation, as per USP specification.

Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts.

Endotoxin Content

NMT 1EU/ml

Storage and Shelf Life:

Store at 2-8°C away from bright light.

Shelf life is 18 months.

Use before expiry date given on the product label.

Disclaimer :

Revision : 04/2022

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.