

MCDB 302 Medium

With Trace elements and L- Glutamine
Without Nucleosides and Sodium bicarbonate

Product Code: AT137

Product Description :

MCDB media were developed for the culture of specific cell types without a serum supplement. The media were supplemented with growth factors, hormones, trace elements, or low levels of dialyzed fetal bovine serum protein (FBSP). Each MCDB medium was formulated for a specific cell type. MCDB 105 and 110 were formulated for rapid clonal growth of normal human diploid cells. MCDB 131 medium was originally developed for the clonal growth of human micro-vascular endothelial cells (HMVEC). MCDB 151, 201 and 302 were originally developed for human keratinocytes, clonal growth of chick embryo fibroblasts and CHO cells.

AT137 is MCDB 302 with trace elements and L-glutamine. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

Composition :

Ingredients	mg/L
INORGANIC SALTS	
Ammonium metavanadate	0.00117
Calcium chloride dihydrate	88.210
Cupric sulphate pentahydrate	0.0025
Disodium hydrogen phosphate anhydrous	141.980
Ferrous sulphate heptahydrate	0.834
Magnesium chloride hexahydrate	122.000
Manganese sulphate	0.000151
Molybdc acid ammonium tetrahydrate	0.0124
Potassium chloride	223.650
Sodium chloride	7599.000
Sodium selenite	0.00173
Zinc sulphate heptahydrate	0.863
AMINO ACIDS	
Glycine	7.510
L-Alanine	8.910
L-Arginine hydrochloride	210.700
L-Asparagine monohydrate	15.000

L-Aspartic acid	13.310
L-Cysteine hydrochloride monohydrate	17.560
L-Glutamic acid	14.710
L-Glutamine	438.600
L-Histidine hydrochloride monohydrate	20.970
L-Isoleucine	3.940
L-Leucine	13.120
L-Lysine hydrochloride	36.540
L-Methionine	4.480
L-Phenylalanine	4.960
L-Proline	34.530
L-Serine	10.510
L-Threonine	11.910
L-Tryptophan	2.040
L-Tyrosine disodium salt dihydrate	7.896
L-Valine	11.720
VITAMINS	
Choline chloride	13.960
D-Biotin	0.00733
D-Ca-Pantothenate	0.238
Folic acid	1.324
Niacinamide	0.0366
Pyridoxine hydrochloride	0.0617
Riboflavin	0.0376
Thiamine hydrochloride	0.337
Vitamin B12	0.136
myo-Inositol	18.020
OTHERS	
D-Glucose	1801.600
Hypoxanthine	4.083
Linoleic acid	0.0841
Phenol red sodium salt	1.242
Putrescine dihydrochloride	0.161
Sodium pyruvate	110.000
Thioctic acid	0.206

Directions :

1. Suspend 11.0gms in 900 ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water.

2. Add 1.18gms of sodium bicarbonate powder (RM447) or 15.73 ml of 7.5% sodium bicarbonate solution (TCL013) for 1 litre of medium and stir until dissolved.
3. Adjust the pH to 0.2 - 0.3 pH units below the desired pH using 1N HCl or 1N NaOH since the pH tends to rise during filtration.
4. Make up the final volume to 1000ml with tissue culture grade water.
5. Sterilize the medium immediately by filtering through a sterile membrane filter with a porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide.
6. Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers.
7. Store liquid medium at 2-8°C and in dark till use.

Material required but not provided :

Tissue culture grade water (TCL010)
Sodium bicarbonate powder (TC230)
Sodium bicarbonate solution, 7.5% (TCL013)
1N Hydrochloric acid (TCL003)
1N Sodium hydroxide (TCL002)
Foetal bovine serum (RM1112/RM10432)

Quality Control:

Appearance

Off-white to Creamish white, homogenous powder.

Solubility

Clear solution at 11.0 gms/L.

pH without Sodium Bicarbonate

6.50 -7.10

pH with Sodium Bicarbonate

7.40 -8.10

Osmolality without Sodium Bicarbonate

260.00 -300.00

Osmolality with Sodium Bicarbonate

280.00 -320.00

Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing it with a control medium through minimum three subcultures.

Endotoxin Content

NMT 5EU/ml

Storage and Shelf Life:

1. All the powdered media and prepared liquid culture media should be stored at 2-8°C. Use before the expiry date. In spite of above recommended storage condition, certain powdered medium may show some signs of deterioration /degradation in certain instances. This can be indicated by change in colour, change in appearance and presence of particulate matter and haziness after dissolution.
2. Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.
3. pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culture vessel used (surface to volume ratio). For example, in large bottles, such as Roux bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH, sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1N HCl or 1N NaOH or by bubbling in carbon dioxide.
4. If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

Revision : 1 / 2011

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

