

Iscove's Modified Dulbecco's Medium(IMDM)

**With L-Glutamine and 25mM HEPES Buffer
Without Glucose and Sodium bicarbonate**

Product Code: AT230

Product Description:

Iscove's Modified Dulbecco's Medium is an enriched modification of Dulbecco's Modified Eagles Medium wherein serum can be partially or totally replaced by chemically defined substances. The medium contains additional amino acids, sodium selenite, sodium pyruvate, vitamins and inorganic salts. Potassium nitrate is substituted by ferric nitrate. IMDM was the first medium utilizing HEPES buffer. The medium when appropriately supplemented supports good growth of precursor cells of erythrocytes and macrophages. The medium also supports good growth of T and B-lymphocytes and a variety of hybrid cells under serum free or reduced serum conditions.

AT230 is Iscove's Modified Dulbecco's Medium with L-glutamine and 25mM HEPES buffer. HEPES, a zwitterionic buffer having a pKa of 7.3 at 37°C prevents the initial rise in pH that tends to occur at the initiation of a culture and increases the buffering capacity of the medium.

Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

Composition:

Ingredients	mg/L
INORGANIC SALTS	
Calcium chloride dihydrate	219.000
Magnesium sulphate anhydrous	97.720
Potassium chloride	330.000
Potassium nitrate	0.076
Sodium chloride	4505.000
Sodium dihydrogen phosphate, anhydrous	109.000
Sodium selenite	0.0173
AMINO ACIDS	
Glycine	30.000
L-Alanine	25.000
L-Arginine hydrochloride	84.000
L-Asparagine monohydrate	25.000

L-Aspartic acid	30.000
L-Cystine dihydrochloride	91.240
L-Glutamic acid	75.000
L-Glutamine	584.000
L-Histidine hydrochloride monohydrate	42.000
L-Isoleucine	104.800
L-Leucine	104.800
L-Lysine hydrochloride	146.200
L-Methionine	30.000
L-Phenylalanine	66.000
L-Proline	40.000
L-Serine	42.000
L-Threonine	95.200
L-Tryptophan	16.000
L-Tyrosine disodium salt dihydrate	104.200
L-Valine	93.600
VITAMINS	
Choline chloride	4.000
D-Biotin	0.013
D-Ca-Pantothenate	4.000
Folic acid	4.000
Nicotinamide	4.000
Pyridoxal hydrochloride	4.000
Riboflavin	0.400
Thiamine hydrochloride	4.000
Vitamin B12	0.013
i-Inositol	7.200
OTHERS	
HEPES Buffer	5958.000
Phenol red sodium salt	15.000
Sodium pyruvate	110.000

Directions:

1. Suspend 13.2gms in 900 ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water.
2. Add 3.024gms of sodium bicarbonate powder (TC230) or 40.32 ml of 7.5% sodium bicarbonate solution (TCL013) for 1 litre of medium and stir until dissolved.
3. Add 4.5gms of Glucose in tissue culture grade water.

4. Adjust the pH to 0.2-0.3 pH units below the desired pH using 1N HCl or 1N NaOH since the pH tends to rise during filtration.
5. Make up the final volume to 1000ml with tissue culture grade water.
6. Sterilize the medium immediately by filtering through a sterile membrane filter with a porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide.
7. Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers.
8. Store liquid medium at 2-8°C and in dark till use.

Material required but not provided:

Tissue culture grade water (TCL010)
Sodium bicarbonate (TC230)
Sodium bicarbonate solution, 7.5% (TCL013)
1N Hydrochloric acid (TCL003)
1N Sodium hydroxide (TCL002)
Fetal bovine serum (RM1112/RM10432)

Quality Control:

Appearance

White to light pink, homogenous powder.

Solubility

Clear solution at 13.2 gms/L.

pH without Sodium Bicarbonate

5.20 - 5.80

pH with Sodium Bicarbonate

6.70 - 7.30

Osmolality without Sodium Bicarbonate(mOsm/Kg H₂O)

210.00 - 250.00

Osmolality with Sodium Bicarbonate(mOsm/Kg H₂O)

280.00 - 320.00

Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts.

Endotoxin content

NMT 1EU/ml

Storage and Shelf Life:

1. All the powdered media and prepared liquid culture media should be stored at 2-8°C. Use before the expiry date. In spite of above recommended storage condition, certain powdered medium may show some signs of deterioration /degradation in certain instances. This can be indicated by change in colour, change in appearance and presence of particulate matter and haziness after dissolution.
2. Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.
3. pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culture vessel used (surface to volume ratio). For example, in large bottles, such as Roux bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH, sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1N HCl or 1N NaOH or by bubbling in carbon dioxide.
4. If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

Revision : 03/2025