



## Dulbecco's Modified Eagle Medium(DMEM), Low glucose

With 1gm Glucose per litre, L-Glutamine and Sodium pyruvate  
Without Calcium chloride and Sodium bicarbonate

**Product Code: AT235**

### Product Description :

Dulbecco's Modified Eagle Medium (DMEM) is one of the most widely used modification of Eagle's medium. DMEM is a modification of Basal Medium Eagle (BME) that contains four fold concentration of amino acids and vitamins. Additionally, the formulation also includes glycine, serine and ferric nitrate. The original formulation contains 1000mg/L of Glucose and was originally used to culture embryonic mouse cells.

AT235 is Dulbecco's Modified Eagle Medium with 1gm per litre of glucose, L-glutamine and sodium pyruvate. It does not contain calcium chloride and sodium bicarbonate.

Users are advised to review the literature for recommendations regarding medium supplementation and Physiological growth requirements specific for different cell lines.

### Composition:

Ingredients	mg/L
<b>INORGANIC SALTS</b>	
Ferric nitrate nonahydrate	0.100
Magnesium sulphate anhydrous	97.720
Potassium chloride	400.000
Sodium chloride	6400.000
Sodium dihydrogen phosphate anhydrous	109.000
<b>AMINO ACIDS</b>	
Glycine	
L-Arginine hydrochloride	30.000
L-Cystine dihydrochloride	84.000
L-Glutamine	62.570
L-Histidine hydrochloride monohydrate	584.000
L-Isoleucine	42.000
L-Leucine	105.000
L-Lysine hydrochloride	105.000
L-Methionine	146.000
L-Methionine	30.000
L-Phenylalanine	66.000
L-Serine	42.000

L-Threonine	95.000
L-Tryptophan	16.000
L-Tyrosine disodium salt dihydrate	103.790
L-Valine	94.000
<b>VITAMINS</b>	
Choline chloride	
D-Ca-Pantothenate	4.000
Folic acid	4.000
Nicotinamide	4.000
Pyridoxal hydrochloride	4.000
Riboflavin	4.000
Thiamine hydrochloride	0.400
i-Inositol	4.000
<b>OTHERS</b>	7.200
D-Glucose	1000.000
Phenol red sodium salt	15.900
Sodium pyruvate	110.000

### Directions :

- Suspend 9.8gms in 900 ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water.
- Add 3.7gms of sodium bicarbonate powder (TC230) or 49.3ml of 7.5% sodium bicarbonate solution (TCL013) for 1 litre of medium and stir until dissolved.
- Adjust the pH to 0.2-0.3 pH units below the desired pH using 1N HCl or 1N NaOH since the pH tends to rise during filtration.
- Make up the final volume to 1000ml with tissue culture grade water.
- Sterilize the medium immediately by filtering through a sterile membrane filter with porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide.
- Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers.
- Store liquid medium at 2-8°C and in dark till use.

## Material required but not provided :

Tissue culture grade water (TCL010)  
Sodium bicarbonate (TC230)  
Sodium bicarbonate solution, 7.5% (TCL013)  
1N Hydrochloric acid (TCL003)  
1N Sodium hydroxide (TCL002)  
Foetal bovine serum (RM1112/RM10434)

## Quality Control:

### Appearance

White to light pink, homogenous powder

### Solubility

Clear solution at 9.8g/L.

### pH without Sodium Bicarbonate

5.90 -6.50

### pH with Sodium Bicarbonate

7.40 -8.00

### Osmolality without Sodium Bicarbonate (mOsm/Kg H<sub>2</sub>O)

230.00 -270.00

### Osmolality with Sodium Bicarbonate (mOsm/Kg H<sub>2</sub>O)

310.00 -350.00

### Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts.

### Endotoxin Content

NMT 1EU/ml

## Storage and Shelf Life:

1. All the powdered media and prepared liquid culture media should be stored at 2-8°C. Use before the expiry date. In spite of above recommended storage condition, certain powdered medium may show some signs of deterioration /degradation in certain instances. This can be indicated by change in colour, change in appearance and presence of particulate matter and haziness after dissolution.
2. Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.

3. pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culture vessel used (surface to volume ratio). For example, in large bottles, such as Roux bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH, sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1N HCl or 1N NaOH or by bubbling in carbon dioxide.
4. If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

## Disclaimer :

Revision : 04/ 2022

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic , research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.