

Technical Data

Blaser-Wang Selective Supplement

FD006

An antibiotic supplement recommended for the selective isolation of *Campylobacter* species.

Composition

Per vial sufficient for 500 ml medium

*Ingredients	Concentration
Polymyxin B	1.250IU
Vancomycin	5mg
Trimethoprim	2.500mg
Amphotericin B	1mg
Cephalothin	7.500mg

Directions:

Rehydrate the contents of 1 vial aseptically with 5 ml sterile distilled water and mix well to dissolve. Avoid frothing of the solution. Aseptically add the contents to 500 ml of sterile, molten, cooled (45-50°C) Brucella Agar Base M074/Brucella HiVegTM Agar Base MV074/Brucella Agar Base, Granulated GM074 /Brucella Broth Base M348 /Brucella HiVegTM Broth Base MV348 /Columbia Blood Agar Base M144/M144A/Columbia Blood Agar Base, HiVegTM MV144/MV144A/ Columbia Blood Agar Base, Granulated GM144/ Columbia Blood HiCynthTM Agar Base MCD144/Columbia Blood HiCynthTM Agar Base W/1% Agar MCD144A. Blood Agar Base No. 2 M834 / M834A / Blood Agar Base No. 2, HiVegTM MV834/ MV834A/ Blood HiCynthTM Agar Base No.2 MCD834 / Blood Agar Base No. 2 w/1.2%Agar, Granulated GM834A with 10% v/ Minerals Growth Supplement Campylobacter FD009. Campylo Thioglycollate Medium Base M908 / Campylo Thioglycollate HiVegTM Medium Base MV908. Campylobacter Agar Base M994/Campylobacter HiVegTM Agar Base MV994 / Christopher's Semisolid Brucella Medium Base M943 or 445 ml of Modified Brucella Agar Base M1268. Mix gently and pour into sterile petri plates.

Type of specimen

Clinical samples: respiratory exudates, faeces etc.; Food samples

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (1,2). For Food samples follow appropriate techniques for handling specimens as per established guidelines (3). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning & Precautions

In Vitro diagnostic use. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Storage and Shelf Life

Store at 2 - 8°C. Use before expiry date on the label.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (1,2).

Reference

- 1. Isenberg (Ed.),2004, Clinical Microbiology Procedures Handbook, Vol.3, American Society for Microbiology, Washington. D.C.
- 2. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
- 3. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, American Public Health Association, Washington, D.C.

* Not For Medicinal Use Revision : 03/2022

HiMedia Laboratories Technical Data



EC REP

HiMedia Laboratories Pvt. Limited, Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) -400604, MS,

CEpartner4U, Esdoornlaan 13, 3951DB Maarn, NL www.cepartner4u.eu



In vitro diagnostic medical device



Storage temperature





Do not use if package is damaged

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.