



# Technical Data

## FO Growth Supplement

FD022

An enrichment supplement recommended for isolation of *Neisseria*.

### Composition

Per bottle

#### Ingredients

Haemoglobin powder

#### Concentration

50G / 100G

### Directions:

A specially prepared powder whose 2% w/v aqueous solution is autoclavable. The aqueous solution is chocolate brown, opaque and contains flocculent dispersible precipitate. It is used for 500 ml medium preparation of GC Agar Base [M434](#) - 5 gms / GC HiVeg™ Agar Base [MV434](#) - 5 gms / Thayer Martin Medium Base [M413](#) - 5 gms / Thayer Martin HiVeg™ Medium Base [MV413](#) - 5 gms / Chocolate No. 2 Agar Base [M1548](#) - 5 gms / Chocolate No. 2 HiVeg™ Agar Base [MV1548](#) - 5 gms / Tellurite Blood Agar Base [M1260](#) - 10 gms / Chocolate Agar Base [M103](#) - 10 gms / Chocolate HiVeg™ Agar Base [MV103](#) - 10 gms / Modified Protease Agar [M1606](#) - 10 gms / Transgrow Medium Base [M1149](#) - 2 gms.

### Type of specimen

Clinical samples - Stool, urine, respiratory exudates, etc.

### Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (1,2). After use, contaminated materials must be sterilized by autoclaving before discarding.

### Warning & Precautions

In Vitro diagnostic use only. For professional use only. Read the label before opening the container. Wear protective gloves/ protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

### Storage and Shelf Life

Store at 2 - 8°C. Use before expiry date on the label.

### Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (1,2).

### Reference

1. Isenberg (Ed.), 2004, Clinical Microbiology Procedures Handbook, Vol.3, American Society for Microbiology, Washington. D.C.
2. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

#### \* Not For Medicinal Use



HiMedia Laboratories Pvt. Limited, Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) -400604, MS, India



CEpartner4U, Esdoornlaan 13, 3951DB Maarn, NL  
[www.cepartner4u.eu](http://www.cepartner4u.eu)



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged

Revision : 03 / 2022

#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.