



Technical Data

V.C.N. Supplement

FD023

An antibiotic supplement, recommended for the selective isolation of *Neisseria gonorrhoeae* and *Neisseria meningitidis*.

Composition

Per vial sufficient for 500 ml medium

*Ingredients

Vancomycin

Colistin methane sulphonate

Nystatin

Concentration

1.500mg

3.750mg

6250Units

Directions:

Rehydrate the contents of 1 vial aseptically with 2 ml sterile distilled water. Mix well and aseptically add it to Thayer Martin Medium Base [M413](#) / Thayer Martin HiVeg™ Medium Base [MV413](#)- for 440 ml of medium aseptically add 50ml sterile lysed blood and one vial of V.C.N. Supplement [FD023](#) along with one vial of Vitamino Growth Supplement [FD025](#). FO Growth Supplement (250ml) [FD022](#) can be used instead of sterile lysed blood in 250ml of medium. In GC Agar Base [M434](#)/ GC HiVeg™ Agar Base [MV434](#) for 250 ml of can be used instead of sterile lysed blood in 250 ml of FO Growth Supplement [FD022](#) and GC Selective Supplement [FD021](#), one vial of GC Selective Supplement [FD021](#) for additional selectivity. If desired V.C.N. Supplement [FD023](#) can be used along with GC Selective Supplement [FD021](#) for additional selectivity.

In Transgrow Medium Base [M1149](#) for 440 ml of medium aseptically add 50 ml of sterile FO Growth Supplement [FD022](#) and one vial of V.C.N. Supplement [FD023](#) along with one vial of Vitamino Growth Supplement [FD025](#).

Type of specimen

Clinical samples - Stool, urine, respiratory exudates, etc.

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (1,2). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning & Precautions

In Vitro diagnostic use only. For professional use only. Read the label before opening the container. Wear protective gloves/ protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Storage and Shelf Life

Store at 2-8°C. Use before expiry date on the label.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (1,2).

Reference

- 1.Isenberg (Ed.),2004, Clinical Microbiology Procedures Handbook, Vol.3, American Society for Microbiology, Washington. D.C.
- 2.Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology,11th Edition. Vol. 1.

Revision : 03/2022

* Not For Medicinal Use



HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India




CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



**In vitro diagnostic
medical device**



CE Marking



Storage temperature



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.