



Sterile BLM 60-600 (Ready to use)

FD364

An innovative enzyme based product that can efficiently inactivate wide range of antibiotics like Penicillins, Cephalosporins of first, second, third & fourth generations and Penems.

Composition

Each vial contains 3 ml of -

*Ingredients

| | |
|--|--------|
| Cephalosporinase Specific activity (IU/vial) | 60 IU |
| Penicillinase Specific activity (IU/vial) | 600 IU |

Note:

1IU is defined as the amount of enzyme needed to hydrolyze 1 μ mole of Penicillin G (Penicillinase) or 1 μ mole of Cephalosporin C (Cephalosporinase) per minute at 25°C and pH 7.0.

1IU of Penicillinase corresponds to 600 Levy Units or 75 Pollock Units.

Directions:

Sterile Beta Lactamase liquid mixture is an optimized ready to use solution that can be directly added to the test samples. The amount of product to be added to the test sample should be determined and set-up depending on the application, concentration of antibiotic to be inactivated, and depending on the specific beta-lactam that should be inactivated.

Aseptic techniques are to be followed throughout the procedure.

Type of specimen

Environmental monitoring

Specimen Collection and Handling

For Environmental samples follow appropriate techniques for handling specimens as per established guidelines (1). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning & Precautions

For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Storage and Shelf Life

Store at 2 - 8°C. Use before expiry date on the label.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (1,2).

Reference


1. Isenberg (Ed.),2004, Clinical Microbiology Procedures Handbook, Vol.3, American Society for Microbiology, Washington. D.C.
2. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

* Not For Medicinal Use

Revision : 00/2022



HiMedia Laboratories Pvt.
Limited, Plot No.C-40, Road
No.21Y, MIDC, Wagle Industrial
Area, Thane (W) -400604, MS,
India




CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



*In vitro diagnostic
medical device*



CE Marking



Storage temperature



Do not use if
package is damaged

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.