

Dextrose Tryptone Agar, Granulated

GM092

Dextrose Tryptone Agar, granulated is recommended for the detection and enumeration of mesophilic and thermophilic aerobic microorganisms in foods.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Dextrose	5.000
Bromocresol purple	0.040
Agar	15.000
Final pH (at 25°C)	6.7±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 30.04 grams in 1000 ml distilled water. Heat to boiling the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Canned foods are most often prone to flat-sour spoilage due to contamination by either mesophilic or thermophilic aerobic spore-formers. Inadequate heat processing is commonly responsible for flat-sour spoilage since spores of mesophilic bacteria are moderately resistant to moist heat. Also *Bacillus stearothermophilus* is the typical species responsible for this type of spoilage (1, 2). *Bacillus coagulans* (*Bacillus thermoacidurans*, a soil organism) is frequently isolated from flat-sour spoilage of canned tomato and dairy products. In flat-sour spoilage, carbohydrates are fermented with the production of lower fatty acids, which sour the product. The small amount of gas produced does not affect the flat appearance of the ends of container.

Dextrose Tryptone Agar, formulated by Williams is recommended for the detection and enumeration of thermophilic flat sour spoilage organisms (3). It is also recommended for general cultural studies by Cameron (4) and other associations (5-8). Dextrose Tryptone Agar is also useful for enumeration of mesophiles and thermophiles in cereal and cereal products, dehydrated fruits, vegetables and spices (9).

Casein enzymic hydrolysate provides essential nutrients to the organisms. Dextrose serves as an energy source by being the fermentable carbohydrate while bromo cresol purple is a pH indicator. Acid producing organisms produce yellow colonies. The plates should be incubated at 55°C for 48 hours in a humid incubator.

While using the agar media, serially diluted test sample are mixed with the media in sterile Petri dishes. Standard procedures issued by various associations should be followed for testing of samples.

Quality Control

Appearance

Light yellow to greenish yellow coloured granular medium

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Purple coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3% w/v aqueous solution at 25°C. pH : 6.7±0.2

pH

6.50-6.90

Cultural Response

Cultural characteristics observed after an incubation at 54-56°C for 36-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Bacillus brevis</i> ATCC 8246	50-100	good-luxuriant (with or without dextrose fermentation)	50-70%	yellow
<i>Bacillus coagulans</i> ATCC 8038	50-100	good-luxuriant	50-70%	yellow
<i>Bacillus stearothermophilus</i> ATCC 7953	50-100	good-luxuriant	50-70%	yellow

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

- Gordon R. E., Haynes and Pang C. H. N., 1973, The Genus *Bacillus*, Agriculture Handbook No. 407, U.S. Department of Agriculture, Washington, D.C.
- Hersom A. C., and Hulland E. D., 1964, Canned Foods, An Introduction to Their Microbiology, (Baumgartner) 5th Ed. Chemical Publishing Company, Inc. New York, N.Y.
- Williams O. B., 1936, Food Res., 1:217.
- Cameron E. J., 1936, J. Assoc. Official Agr. Chem., 19:433.
- Association of Official Analytical Chemists, 1978, Bacteriological Analytical Manual, 5th Edition, AOAC, Washington, D.C.
- Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed. APHA Inc., Washington, D.C.
- National Canners Association, 1968, Laboratory Manual for Food Caners and Processors, Vol. I
- National Canners Association, 1954, A Laboratory Manual for the Canning Industry, 1st Edition, National Canners Associations, Washington.
- Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.