

Standard Nutrient Agar No. 2, Granulated

GM1627

Standard Nutrient Agar No. 2, granulated is recommended for the cultivation and enrichment of less fastidious bacteria. It can be used in detection of inhibitors in bacteriological examination of meat.

Composition**

Ingredients	Gms / Litre
Meat peptone	3.450
Casein enzymic hydrolysate	3.450
Sodium chloride	5.100
Agar	13.000
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 25 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Fastidious organisms are organisms which require preformed organic molecules like vitamins, amino acids, nucleic acids, carbohydrates. In general bacterial pathogens need more preformed organic molecules than do nonpathogens. Media which are highly nutritional are generally used to enrich less fastidious organism so as to isolate them from test samples. Standard Nutrient Area No.2 can be used in the detection of inhibitors during the bacteriological examination of meat (1). This medium can also be modified with various additives (2). Standard Nutrient Broth No.2 can also be used for the examination of water (3).

Meat Peptone and casein enzymic hydrolysate in the medium provides the nitrogenous and carbon source with other essential nutrients. Sodium chloride maintains the osmotic equilibrium of the medium.

Quality Control

Appearance

Cream to yellow coloured granular medium

Gelling

Firm, comparable with 1.3% Agar gel

Colour and Clarity of prepared medium

Dark amber to amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 2.5% w/v aqueous solution at 25°C. pH : 7.5±0.2

pH

7.30-7.70

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Escherichia coli</i> ATCC 11775	50-100	good-luxuriant	≥70%
<i>Shigella flexneri</i> ATCC 29903	50-100	good-luxuriant	≥70%
<i>Salmonella</i> Typhimurium ATCC 13311	50-100	good-luxuriant	≥70%

<i>Staphylococcus aureus</i> ATCC 6538 P	50-100	good-luxuriant	$\geq 70\%$
<i>Streptococcus pyogenes</i> ATCC 21059	50-100	good-luxuriant	$\geq 70\%$
<i>Listeria monocytogenes</i> ATCC 19118	50-100	good-fair	$\geq 50\%$

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Levetzow,R: Untersuchung auf Hemmstoffe im Rahmen der bakteriologischen Fleischuntersuchung, - Bundesgesundheitsblatt, 1971.14; 211-213.
2. Zavanella, M., Aurelia, P., a. Ferrini, A.M: Improved microbiological method for the detection of antimicrobial residues in meat.- 1986. Arch Lebensmittelhyg.,37:118-120.
3. Din Deutsches Institut fur Normung e.V: Deutsche Einheitsverfahren zur Wasser-, Abwasser- und Schlammuntersuchung, Mikrobiologische Verfahren (Gruppe K). Nachweis von Pseudomonas aeruginosa (K 8). DIN 38411.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.