

## Mueller Kauffman Tetrathionate Broth Base, Granulated

GM876

Mueller Kauffman Tetrathionate Broth, granulated is used for improved enrichment and isolation of Salmonellae.

### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	7.000
Papaic digest of soyabean meal	2.300
Sodium chloride	2.300
Calcium carbonate	25.000
Sodium thiosulphate	40.700
Ox bile	4.750

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 82.05 grams in 1000 ml distilled water. Heat just to boiling. DO NOT AUTOCLAVE. Cool and just before use aseptically add 19 ml of iodine solution (20 g iodine and 25 g potassium iodide in 100 ml sterile distilled water) and 9.5 ml of 0.1% brilliant green solution. Mix well before dispensing in sterile tubes to disperse calcium carbonate uniformly .

Note: Due to presence of calcium carbonate, the prepared media forms opalescent solution with white precipitate

### Principle And Interpretation

The examination of various types of food products for *Salmonella* requires methods different from those used in clinical laboratories. The need for such method is due to the generally low numbers of Salmonellae in foods and the frequently poor physiological state of these pathogens following exposure to stressful conditions during food processing or storage. Injured *Salmonella* are resuscitated in non-selective broth medium, which facilitates detection of sublethally injured *Salmonella* . The ideal pre-enrichment broth should provide for the repair of cell damage, dilute toxic or inhibitory substances and nutritive enough to favour growth of *Salmonella* . In the analysis of food for *Salmonella* , pre-enrichment cultures are usually incubated at 35-37°C for 18-24 hours and then a portion is subcultured to one or more selective enrichment broths. Normally 1 ml of pre-enrichment culture is inoculated to 9 ml of selective enrichment broth. Selective enrichment media contains selective ingredients that allow the proliferation of *Salmonella* and inhibit the growth of competing non-salmonella microorganisms. Lactose Broth (M1003) is recommended by BAM for pre-enrichment of *Salmonella* from food. Selective enrichment is done in Tetrathionate Broth and Rappaport Vassiliadis Medium. For the detection of foodborne *Salmonella* , various modifications of Tetrathionate Broth have generally found wider applications (7).

Mueller (1) recommended Tetrathionate Broth as a selective medium for the isolation of *Salmonella* . Kauffman (2) modified the formula to include ox bile and brilliant green as selective agents to suppress bacteria such as *Proteus* species. The British Standard Specification specifies Brilliant Green Tetrathionate Broth for isolating *Salmonella* from meat and meat products and from poultry and poultry products (3). It is also a recommended selective broth for isolating *Salmonella* from animal feces and sewage-polluted water (4). Selectivity is conferred by tetrathionate (from the reaction of thiosulphate and iodine). Using more than one selective broth increases the isolation of *Salmonella* from samples with multiple serotypes (5). Mueller Kauffman Tetrathionate Broth Base conforms to ISO specifications (9).

Mueller Kauffman Tetrathionate Broth Base contains casein enzymic hydrolysate and papaic digest of soyabean meal as sources of carbon, nitrogen, vitamins and minerals. Ox bile and added brilliant green are selective agents, which inhibit gram-positive and other gram-negative organisms. Calcium carbonate is the buffer. Sodium chloride maintains osmotic equilibrium. Sodium thiosulphate is a source of sulfur. The tetrathionate (S<sub>4</sub>O<sub>6</sub>) anions constitute the principle selective agent in these enrichment media. If desired, 4 mg of novobiocin per litre of broth can be added to suppress *Proteus* species (6).

Add approximately 10 grams of sample to 100 ml of broth. Shake well and place the flask in a 45°C water bath for 15 minutes. Remove the flasks and place in an incubator or water bath at 43°C. Several studies have shown increased recovery of *Salmonella* following incubation of selective enrichment at 43°C (8). After an incubation for 18-24 hours and 48 hours, subculture on Brilliant Green Agar, Modified (M016). This medium is not suitable for the growth of *Salmonella* Typhi, *Salmonella* Sendai, and *Salmonella* Pullorum etc.

The complete medium is unstable and should be used immediately. It may be stored at 2-8°C in the dark for no more than 7 days.

Organisms other than Salmonellae, such as *Morganella morganii* and some *Enterobacteriaceae* may grow in the medium. Therefore, confirmatory tests should be carried out on all presumptive *Salmonella* colonies that are recovered.

## Quality Control

### Appearance

Cream to yellow coloured granular medium

### Colour and Clarity of prepared medium

With added brilliant green and iodine solution - Light green coloured opalescent solution forms with heavy white precipitate

### Cultural Response

Cultural characteristics observed, when subcultured on Soyabean Casein digest Agar, after an incubation at 43°C for 18-24 hours with added iodine and brilliant green solution.

### Cultural Response

Organism	Inoculum (CFU)	Recovery
<b>Cultural Response</b>		
<i>Salmonella</i> Typhimurium ATCC 14028	50-100	excellent
<i>Salmonella</i> Enteritidis ATCC 13076	50-100	excellent
<i>Salmonella</i> Paratyphi A ATCC 9150	50-100	excellent
<i>Salmonella</i> Paratyphi B ATCC 8759	50-100	excellent
<i>Salmonella</i> Typhi ATCC 6539	$\geq 10^3$	inhibited
<i>Escherichia coli</i> ATCC 25922	50-100	none-poor
<i>Proteus vulgaris</i> ATCC 13315	50-100	none-poor
<i>Shigella flexneri</i> ATCC 12022	$\geq 10^3$	inhibited

## Storage and Shelf Life

Store below 30°C in tightly closed container and use the freshly prepared medium. Use before expiry date on the label.

## Reference

- Mueller L., 1923, C. R. Soc. Biol., (Paris) 89:434.
- Kauffman F., 1935, Ztschr. F. Hyg., 117:26.
- International Organization for Standardization, 1974, (Draft International Standard ISO/DIS 3565), Geneva, Switzerland.
- Public Health Laboratory Service, 1974, Monograph Series No. 8, Public Health Laboratory Service, London, England.
- Harvey R. W. S. and Price T. S., 1976, J. Hyg. Camb., 77:333.
- Jeffries L., 1959, J. Clin. Pathol., 12:568.
- Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.
- DAoust J. Y., 1989, Salmonella in Food borne Bacterial pathogens, (Eds.) Doyle M. P., 327, Marcel Dekker, New York.
- International Organization for Standardization (ISO), 2002, Draft 6579.

Revision : 00 / 2014

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.