

M17 Agar Base ,Granulated[®]

GM929

Intended Use:

Recommended for cultivation of lactic Streptococci and plaque assay of lactic bacteriophages.

Composition**

Ingredients	g / L
Peptone	5.000
Soya peptone	5.000
Yeast extract	2.500
HM peptone B #	5.000
Ascorbic acid	0.500
Magnesium sulphate	0.250
Lactose	5.000
Agar	10.000
Final pH (at 25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

-Equivalent to Beef extract

Directions

Suspend 33.25 grams in 1000 ml purified/ distilled water. Add 19 grams of Disodium β-Glycerophosphate. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and dispense as desired.

Principle And Interpretation

M17 media are based on the formulation described by Terzaghi and Sandine (1) for the cultivation and enumeration of lactic Streptococci and their bacteriophages. It is possible to study plaque morphology and lysogeny. M17 Agar is recommended by the International Dairy Federation (2) for selective enumeration of *Streptococcus thermophilus* from yoghurt. M17 Agar is recommended by APHA for the cultivation of lactic Streptococci (3,4). Lactic Streptococci are nutritionally fastidious and require complex media for optimal growth (5,6). Disodium glycerophosphate maintains the pH above 5.7. The maintenance of pH is very important as lower pH results in injury and reduced recovery of lactic Streptococci. Glycerophosphate does not form precipitate with calcium which is needed for the plaque assay of lactic bacteriophages

Peptone, soya peptone, yeast extract and HM peptone B provide carbonaceous, nitrogenous compounds, long chain amino acids, vitamin B complex and other essential growth factors. Lactose is the fermentable carbohydrate and ascorbic acid is stimulatory for the growth of lactic Streptococci. Magnesium sulphate provides essential ions to the organisms. Disodium-#-glycerophosphate maintains the pH above 5.7. The maintenance of pH is very important as lower pH results in injury and reduced recovery of lactic Streptococci. Disodium glycerophosphate suppresses *Lactobacillus bulgaricus*. Shankar and Davies (7) reported isolation and enumeration of *Streptococcus thermophilus* from yoghurt. M17 Agar is also recommended by the International Dairy Federation (2) for selective enumeration of *Streptococcus thermophilus* from yoghurt. It is also suitable for cultivation and maintenance of starter cultures for cheese and yoghurt manufacturing. This medium helps in detecting streptococcal mutants that are lactose non-fermenters. Suggested technique to enumerate streptococci is to seed in mass or by stabbing with agar, melted and cooled to 50-55° C, and incubating them at 42°C for 24 hours period. With these conditions, all the colonies might be streptococci. Longer incubation periods or lower temperatures may cause morphological changes in the colonies, which hinders in the recognition of the colonies. Lactose-positive colonies of streptococci are visible after 15 hours and after 5 days they may reach a diameter of about 3-4 mm, whereas those of lactose-negative are 1 mm in diameter. Bacteriophages presence is observed by appearance of characteristic plaques over the bacterial growth.

Type of specimen

Food and Dairy samples

Specimen Collection and Handling:

Suggested technique to enumerate streptococci is to seed in mass or by stabbing with agar, melted and cooled to 50-55°C, and incubating them at 42°C for 24 hours period. With these conditions, all the colonies might be streptococci. Longer incubation periods or lower temperatures may cause morphological changes in the colonies, which hinders in the recognition of the colonies. Lactose-positive colonies of streptococci are visible after 15 hours and after 5 days they may reach a diameter of about 3-4 mm, whereas those of lactose-negative are 1 mm in diameter. Bacteriophages presence is observed by appearance of characteristic plaques over the bacterial growth.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. The maintenance of pH is very important as lower pH results in injury and reduced recovery of lactic Streptococci.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow granulated free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel.

Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 3.33% w/v aqueous solution at 25°C. pH : 7.1±0.2

pH

6.90-7.30

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours with added Disodium β-Glycerophosphate.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	50-100	good-luxuriant	≥50%
<i>Lactobacillus bulgaricus</i> subsp. <i>bulgaricus</i> ATCC 11842 (00102*)	50-100	none-poor	≤10%
<i>Lactobacillus leichmannii</i> ATCC 4797	50-100	good-luxuriant	≥50%
# <i>Lactiplantibacillus plantarum</i> ATCC 8014	50-100	good-luxuriant	≥50%
<i>Streptococcus thermophilus</i> ATCC 14485	50-100	good-luxuriant	≥50%

Key : *Corresponding WDCM numbers.

#Formerly known as *Lactobacillus plantarum*

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (8,9).

Reference

1. Terzaghi B.E. and Sandine W.E., 1975, Appl. Microbiol., 29:807.
2. International Dairy Federation, 1981, Joint IDF/ISO/AOAC Group E44.
3. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
4. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.
5. Anderson A.W. and Elliker P.R., 1953, J. Dairy Sci., 36:161.
6. Reiter B. and Oran J.D., 1962, J. Dairy Res., 29:63.
7. Shankar P.A. and Davies F.L., 1977, Soc. Dairy Technol., 30:28.
8. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
9. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision :01 / 2025

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.