



Soyabean Casein Digest Medium

LQ027XCK

Intended Use

Sterility test media prepared in accordance with harmonized methods of USP, EP, BP, JP & IP.

Composition**

Ingredients	g / L
Tryptone\$	17.000
Soya peptone^	3.000
Sodium chloride	5.000
Glucose monohydrate	2.500
Dipotassium hydrogen phosphate	2.500
Final pH after sterilization (at 25°C)	7.3±0.2
**Formula adjusted, standardized to suit performance parameters	
\$ Equivalent to Pancreatic digest of casein	^ Equivalent Papaic digest of soyabean meal

Directions

Label the ready to use LQ027XCK bottle. Inoculate the sample and Incubate at specified temperature and time.

Principle And Interpretation

Soyabean Casein Digest Medium is recommended as a sterility testing medium in accordance with the harmonized method of USP/EP/BP/JP/IP (1-5). It is used for the sensitivity testing of antimicrobial agents by the tube dilution method (6). It is also employed in diagnostic research in microbiology. This medium is used as a diluent and suspending medium for preparation of samples or test strains. It is also employed in sample preparation for testing of products, wherein incubation is carried out, only to serve sufficient resuscitation of the cell, while avoiding multiplication of the organism.

Tryptone and soya peptone makes this medium nutritious by providing nitrogenous and carbonaceous compounds, long chain peptides, vitamins and other essential nutrients for the growth of microorganisms. Natural sugars in soybean promote growth of fastidious organism. Glucose monohydrate is the fermentable source of carbon and dibasic potassium phosphate serves as the buffer in the medium. Sodium chloride maintains the osmotic balance of the medium. This medium is recommended for sterility checking and for studying total aerobic microbial count in verification of microbiological testing procedures employed for sterility checking.

Type of specimen

Pharmaceutical samples

Specimen Collection and Handling

For pharmaceutical samples, follow appropriate techniques for sample collection, processing as per guidelines (1-5). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Biochemical characterization is necessary to be performed on colonies from pure cultures for further identification.
2. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
3. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Sterile Soyabean Casein Digest medium in a glass bottle.

Colour

Light yellow coloured clear solution

Quantity of Medium

90 ml of medium in glass bottle.

pH

7.10- 7.50

Growth Promotion Test

In accordance with the harmonized method of USP/EP/BP/JP.

Sterility Check

Passes release criteria.

Growth promoting properties

Clearly visible growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating ≤ 100 cfu (at 30-35°C for 18-24 hours for bacteria). Growth promotion is carried out as per USP/EP/BP/JP.

Sterility Testing + Validation

The medium is tested with suitable strains of microorganisms inoculating ≤ 100 cfu and incubating at 20-25°C for not more than 3 days in case of bacteria and not more than 5 days in case of fungi.

Organism	Inoculum (CFU)	Growth ^s	Incubation period	Incubation temperature
Growth promoting				
<i>Salmonella</i> Abony NCTC 6017 (00029*)	≤ 100	luxuriant	18 -24 hrs	30 -35 °C
<i>Salmonella</i> Typhimurium ATCC 14028 (00031*)	≤ 100	luxuriant	18 -24 hrs	30 -35 °C
^ <i>Pseudomonas paraeruginosa</i> ATCC 9027 (00026*)	≤ 100	luxuriant	18 -24 hrs	30 -35 °C
<i>Escherichia coli</i> ATCC 8739 (00012*)	≤ 100	luxuriant	18 -24 hrs	30 -35 °C
\$\$ <i>Bacillus spizizeni</i> ATCC 6633 (00003*)	≤ 100	luxuriant	18 -24 hrs	30 -35 °C
## <i>Kokuria rhizophila</i> ATCC 9341	≤ 100	luxuriant	18 -24 hrs	30 -35 °C
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*)	≤ 100	luxuriant	18 -24 hrs	30 -35 °C
Sterility Testing- Growth promotion+Validation				
^ <i>Pseudomonas paraeruginosa</i> ATCC 9027 (00026*)	≤ 100	luxuriant	≤ 3 d	20 -25 °C
\$\$ <i>Bacillus spizizenii</i> ATCC 6633 (00003*)	≤ 100	luxuriant	≤ 3 d	20 -25 °C
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*)	≤ 100	luxuriant	≤ 3 d	20 -25 °C
## <i>Kokuria rhizophila</i> ATCC 9341	≤ 100	luxuriant	≤ 3 d	20 -25 °C
<i>Candida albicans</i> ATCC 10231 (00054*)	≤ 100	luxuriant	≤ 5 d	20 -25 °C
# <i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	≤ 100	luxuriant	≤ 5 d	20 -25 °C

Key : (*) Corresponding WDCM numbers,

- Formerly known as *Aspergillus niger*,

^ Formerly known as *Pseudomonas aeruginosa*

\$ - Luxuriant growth refers to turbid growth

- Formerly known as *Micrococcus luteus*

\$\$- Formerly known as *Bacillus subtilis* subsp. *spizizenii*

Storage and Shelf Life

Store between 15-30°C. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (7,8).

Reference

- 1.The British Pharmacopoeia, 2022, Medicines and Healthcare products Regulatory Agency.
- 2.European Pharmacopoeia, 2022, 10 th volume, European Directorate for the quality of medicines & Healthcare.
- 3.The Japanese Pharmacopoeia, 17th edition, 2016, The Ministry of Health, Labour and welfare.
- 4.Indian Pharmacopoeia, Addendum 2024, Indian Pharmacopoeia Commission, Ministry of Health and Family Welfare Government of India.
- 5.The United States Pharmacopoeia-National Formulary (USP-NF), 2022.
- 6.Wright and Welch, 1959-60, Antibiotics Ann., 61.
- 7.Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
- 8.Jorgensen,J.H., Pfaller , M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision :01/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

